# Index for Volume 52 (2003)

*Syst. Biol.* 52(6):876–911, 2003
Copyright © Society of Systematic Biologists
ISSN: 1063-5157 print / 1076-836X online
DOI: 10.1080/10635150390276444

COMPILED BY JOE MADIA

12S rDNA, 407, 727, 798–799
12S rRNA, 39, 42, 56–58, 161, 163–165, 167, 181, 206–207, 209–212, 215, 218, 297, 440, 443, 445–447, 451, 456, 460, 462–467, 605
16S mtDNA, 738
16S rDNA, 72–76, 407, 727, 798–799
16S rRNA, 206–207, 209–212, 215, 219, 297–298, 307, 440, 443, 445–447, 451, 456, 460, 462–467, 557, 559, 618–640, 758–759, 762
18S rDNA, 57, 72–75, 820, 823
18S rRNA, 283–295, 296, 298, 307, 311–333, 557, 559
26S rDNA, 820, 823
28S rDNA, 57, 72–75
28S rRNA, 285, 287, 298, 307, 557, 559, 618, 620–623, 745–756

A2AB, 605, 614–615
A6, 674
Abdo, Zaid, *see* Minin, Vladimir, 674–683
*Acacia*
  *koa*, 820
*Acaenosuchus*, 241–243, 250–251
*Acanthalictus*, 24, 26, 32
Acari, 298, 430
Acaridae, 430
*accD*, 336
Accelerated likelihood surface exploration: the likelihood ratchet, R. A. Vos, 368–373
Accuracy, 55–65, 428–435
*Achalinus*, 451
  *rufescens*, 440
Achramorpha, 314–315, 328, 333
*Acleisanthes*, 334–351
  *acutifolia*, 341, 343
  *angustifolia*, 341–343, 349
  *chenopodioides*, 341
  *crassifolia*, 347
  *lanceolata*, 341, 348
  *longiflora*, 347
  *parvifolia*, 341, 347–348
  *purpusiana*, 341
  *undulata*, 341, 343, 349
  *wrightii*, 343, 347
*Aclys*, 758, 775
  *concinna*, 758
*Acomastylis*, 374
*Acrocera bulla*, 756
Acroceridae, 756
Acrochordidae, 439–440, 442, 447, 455, 457
*Acrochordus*, 439, 442–443, 451–452, 457
  *granulatus*, 440
  *javanicus*, 440
Adams consensus tree, 850
Adams, Dean C., *see* Naylor, Gavin J. P., 864–865
Adams, Richard J., *see* Johnson, Kevin P., 37–47
Adams tree, 463
Adaptation, 501–514
Adaptive radiation, 70, 89–92, 207
Advanced snakes, 439–459
*Aepyceros*, 209, 212, 215
  *melampus*, 210
Aepycerotini, 209–210
Aetosaur, 253–255
  phylogeny, 852
Aetosauria, 239
*Aetosauroides*, 241, 243, 248, 250–252
  *scagliai*, 250
*Aetosaurus*, 241–243, 248, 250–252
Affine, 66–69
African elephants, 591
Agamidae, 757
Agamid lizards, 758
Agaricales, 430
*Agkistrodon*, 447, 451
  *contortrix*, 440
*Agrionemis exsudans*, 94
Agriocnemus, 94
*Ailurus fulgens*, 846
Akaike Information Criterion (AIC), 79, 89, 95, 824
Alethinophidia, 440
*Alexeter* sp., Alcini, 209
Algae, 430, 518, 524
Algorithm, 696–704
Alinkidae, 751
Alivon, Eliane, *see* Manuel, Michael, 311–333
Alcelaphini, 209–210
Alleles, 374
  mitochondrial, 41
  nuclear, 41, 103
*Alleno pithecus nigroviridis*, 736
Alligator, American, 388
*Alligator*, 391, 409
  *mississippiensis*, 286–287, 386, 388, 391, 405
  *sinensis*, 405
Alligatoridae, 394, 396
Alligatoridea, 403, 405, 408–409, 412, 415
Allodape, 5, 7–8
  mucronata, 2
  skaifeorum, 2
Allodapine, 2, 7–9, 11
  bees, 1
Allodapini, 1–14
*Allodapula*, 2, 9–10, 12
*Allodapulodes*, 9
Allopolyploidy, 374–385
Allothyrus sp., 300, 306
Allozyme, 23, 504, 506, 508–509
  trees, 509
α-lactalbumin, 206, 209–212, 215, 218–220

# Index for Volume 52 (2003)

*Syst. Biol.* 52(6):876–911, 2003
Copyright © Society of Systematic Biologists
ISSN: 1063-5157 print / 1076-836X online
DOI: 10.1080/10635150390276444

COMPILED BY JOE MADIA

12S rDNA, 407, 727, 798–799
12S rRNA, 39, 42, 56–58, 161, 163–165, 167, 181, 206–207, 209–212, 215, 218, 297, 440, 443, 445–447, 451, 456, 460, 462–467, 605
16S mtDNA, 738
16S rDNA, 72–76, 407, 727, 798–799
16S rRNA, 206–207, 209–212, 215, 219, 297–298, 307, 440, 443, 445–447, 451, 456, 460, 462–467, 557, 559, 618–640, 758–759, 762
18S rDNA, 57, 72–75, 820, 823
18S rRNA, 283–295, 296, 298, 307, 311–333, 557, 559
26S rDNA, 820, 823
28S rDNA, 57, 72–75
28S rRNA, 285, 287, 298, 307, 557, 559, 618, 620–623, 745–756

A2AB, 605, 614–615
A6, 674
Abdo, Zaid, *see* Minin, Vladimir, 674–683
*Acacia*
  *koa*, 820
*Acaenosuchus*, 241–243, 250–251
*Acanthalictus*, 24, 26, 32
Acari, 298, 430
Acaridae, 430
*accD*, 336
Accelerated likelihood surface exploration: the likelihood ratchet, R. A. Vos, 368–373
Accuracy, 55–65, 428–435
*Achalinus*, 451
  *rufescens*, 440
Achramorpha, 314–315, 328, 333
*Acleisanthes*, 334–351
  *acutifolia*, 341, 343
  *angustifolia*, 341–343, 349
  *chenopodioides*, 341
  *crassifolia*, 347
  *lanceolata*, 341, 348
  *longiflora*, 347
  *parvifolia*, 341, 347–348
  *purpusiana*, 341
  *undulata*, 341, 343, 349
  *wrightii*, 343, 347
*Aclys*, 758, 775
  *concinna*, 758
*Acomastylis*, 374
*Acrocera bulla*, 756
Acroceridae, 756
Acrochordidae, 439–440, 442, 447, 455, 457
*Acrochordus*, 439, 442–443, 451–452, 457
  *granulatus*, 440
  *javanicus*, 440
Adams consensus tree, 850
Adams, Dean C., *see* Naylor, Gavin J. P., 864–865
Adams, Richard J., *see* Johnson, Kevin P., 37–47
Adams tree, 463
Adaptation, 501–514
Adaptive radiation, 70, 89–92, 207
Advanced snakes, 439–459
*Aepyceros*, 209, 212, 215
  *melampus*, 210
Aepycerotini, 209–210
Aetosaur, 253–255
  phylogeny, 852
Aetosauria, 239
*Aetosauroides*, 241, 243, 248, 250–252
  *scagliai*, 250
*Aetosaurus*, 241–243, 248, 250–252
Affine, 66–69
African elephants, 591
Agamidae, 757
Agamid lizards, 758
Agaricales, 430
*Agkistrodon*, 447, 451
  *contortrix*, 440
*Agrionemis exsudans*, 94
Agriocnemus, 94
*Ailurus fulgens*, 846
Akaike Information Criterion (AIC), 79, 89, 95, 824
Alethinophidia, 440
*Alexeter* sp., Alcini, 209
Algae, 430, 518, 524
Algorithm, 696–704
Alinkidae, 751
Alivon, Eliane, *see* Manuel, Michael, 311–333
Alcelaphini, 209–210
Alleles, 374
  mitochondrial, 41
  nuclear, 41, 103
*Alleno pithecus nigroviridis*, 736
Alligator, American, 388
*Alligator*, 391, 409
  *mississippiensis*, 286–287, 386, 388, 391, 405
  *sinensis*, 405
Alligatoridae, 394, 396
Alligatoridea, 403, 405, 408–409, 412, 415
Allodape, 5, 7–8
  mucronata, 2
  skaifeorum, 2
Allodapine, 2, 7–9, 11
  bees, 1
Allodapini, 1–14
*Allodapula*, 2, 9–10, 12
*Allodapulodes*, 9
Allopolyploidy, 374–385
Allothyrus sp., 300, 306
Allozyme, 23, 504, 506, 508–509
  trees, 509
α-lactalbumin, 206, 209–212, 215, 218–220

*Alsinidendron*, 832
Alsinoideae, 833
*Alsophis*, 447, 451
*portoricensis*, 440
Amato, George, *see* Gatesy, John, 403–422
Amber fossils, 746
*Amblonyx*, 590
*cinereus*, 573, 577
*Amblyomma*, 299, 304, 309–310
*americanum*, 299
*cajennense*, 299
*fimbriatum*, 299
*glebopalma*, 299
*hebraeum*, 299, 306
*latum*, 299
*maculatum*, 299
s.l., 310
*varanensis*, 299
*variegatum*, 299
*vikirri*, 299
Amblyomminae, 299
Amblypygi, 62
*Ambystoma mexicanum*, 287
American alligator, 388
Amino acids, 3, 6, 10, 95, 165, 296, 585, 594, 596, 605, 662
*Ammocrypta*, 491, 493, 496–499
*beanii*, 491, 497, 499
*vivax*, 491, 496–497, 499
Among-site rate variation, 604–617
Amphibians, 286, 524
Amphimerycidae, 222
Amphimeryx, 222
Amphoriscidae, 314
Amphoriscus, 314–315, 332
Analysis,
Bayesian, 26, 30, 89, 96, 102, 133, 140, 152, 165, 167, 181–182, 211, 390, 439–459, 477–487, 571, 576, 583, 604–617, 618, 621, 623, 629, 665, 681, 708, 713–714, 760, 797, 801–803, 807
Bayesian information, 377
biogeographic, 258, 761, 824
biotic element, 717–723
bootstrap, 7, 26, 447, 580, 609, 611, 621, 697, 775, 824, 841
Bremer support, 211
canonical correspondence, 687–689, 693
canonical variate, 190, 196, 198, 491, 496
cladistic, 9, 81, 253, 403, 541, 572, 738
cluster, 194, 205, 497, 722
continuous jackknife function (CJF), 55–58, 61–63
copath, 41
cophylogenetic, 737
correspondence, 687
cytogenetic, 374
decay (*see also* Bremer support), 128, 243, 407
distance, 334, 343
DNA, 551
generalized Procrustes, 66, 68
genetic, 197
jackknife, 407
Jungles, 39, 43, 45
likelihood, 18, 263, 374, 708, 715, 803–804, 807
local clock, 89, 715
logDet, 3, 6–7, 11
Markov Chain Monte Carlo, 140, 213, 486, 750, 801
maximum likelihood, 1, 3, 7, 11, 26–27, 30, 42, 63, 76, 79, 89, 95–97, 99, 166–167, 171, 182, 211–212, 215, 231, 318, 322, 327, 329, 341, 377–378, 386, 389–90, 394–396, 407, 419, 715, 737, 740, 760, 797, 801, 824, 850
maximum parsimony, 3, 5–7, 10–11, 95, 166–167, 182, 211, 213, 215, 275, 317, 322, 327, 737, 797, 801
molecular, 103–104, 221, 323, 386, 397, 542
molecular dating, 375
morphological, 187, 213, 323, 501, 504, 541, 560, 605, 858
morphometric, 197
multigene, 604
multivariate, 652
neighbor-joining, 463, 530, 532
nested-clade, 159–185
NHML, 390
nonparametric bootstrap, 166
paleontological, 605
parsimony, 7, 26, 30, 42, 45, 70, 76, 79, 83, 249, 251, 255, 257, 260, 262, 271–280, 336, 377, 389–390, 392, 406, 412, 423, 506, 508, 530, 554, 571, 575, 583, 618, 620–621, 623, 629, 727, 764, 802, 804, 824
partitioned, 4, 396, 604–617
partitioned likelihood, 386–402
phylogeographic, 461
principal component, 687
principal coordinate, 860
reconciliation, 39, 42
redundancy, 687
regression, 68
relative warp, 198
reweighted, 318
sensitivity, 554–564
sequence, 737
SSR, 612
stability, 554–564
star topology, 481
statistical, 131, 847
supermatrix, 724, 726, 728, 853
supertree, 725–726, 728, 853
total evidence, 839
tree-reconstruction, 182
UPGMA cluster, 488, 491, 493, 496–497
variable rate, 751
*Anamixilla*, 314–315, 320, 322–324
*torresi*, 316
*Anas platyrhynchos*, 286
Anatomical characters, 742
Ancestral character state reconstruction, 820–841
Ancient Allopolyploid speciation in Geinae (Rosaceae): evidence from nuclear granule–bound starch synthase (GBSSI) gene sequences, Jenny E. E. Smedmark, Torsten Eriksson, Rodger C. Evans, and Christopher S. Campbell, 374–385
Andayani, Noviar, *see* Evans, Ben J., 794–823
*Anemonia sulcata*, 316
Angiosperms, 346, 348, 518, 524, 753, 784
Animals, 89, 591, 736, 742, 820
Anisopodidae, 756
*Anocentor*, 305
*Anolis*, 423, 105
Anomalodesmata, 557
Anomaluridae, 605–606, 614
Anomaluroidea, 614–615
Anomaluromorpha, 605–606
*Anonyx*, 590
*capensis*, 573–574, 576–577
*Anopheles*, 746, 754
*gambiae*, 745
*Anseranas semipalmata*, 388
*Antedon*, 317
*serrata*, 316
Antelope, 206
Anthophyte, 787–788

Anthropoids, 48, 53, 709, 711–713
*Antilocapra*, 207, 209, 212–213, 215, 218, 220–221
  *americana*, 210, 219, 224
Antilocapridae, 206–207, 209–210, 212–213, 218–219, 224
Antilocaprinae, 210, 218
Antilopini, 209–210
*Antipathes lata*, 316
Ants, 23, 33
*Aotus* 740–742
  *nancymai*, 736, 738, 742
Aparallactinae, 447
Aparallactus, 446, 451, 455–456
  *guentheri*, 440, 459
  *werneri*, 440
Apes, 524, 707–708
Aphids, 23, 137, 139–140, 143, 146
*Aphroceras*, 314–315, 332
Aphyllophorales, 430
Apidae, 1
*Apiocera haruspex*, 756
Apioceridae, 746, 750
*Apis mellifera*, 26, 28, 296
Aplocheiloid fishes, 549–553
Aplocheilidae, 548, 552
Aplocheiloidei, 552
*Aplodontia rufa*, 606
Aplodontidae, 605–606, 608, 614
Aplopappus, 33
*Aplopeltura*, 451
  *boa*, 440, 446
*Aplysilla sulfurea*, 316
Approximately unbiased test, 781–793
*Aprasia*, 757–758, 762, 764, 767, 770–771, 773, 776–777
  *aurita*, 759
  *fusca*, 759, 776
  *inaurita*, 759
  *parapulchella*, 759, 776
  *picturata*, 759
  *pseudopulchella*, 759, 776
  *pulchella*, 759, 776
  *repens*, 759, 776
  *rostrata*, 758, 776
  *smithi*, 759
  *striolata*, 759, 770
Apterygota, 328
*Apteryx*
  *australis*, 286, 471–472, 474
  *mantelli*, 471, 474
*Apus affinus*, 286
Arachnids, 56, 61, 299
Araneae, 62, 70, 300, 557
Araneid, 56, 61
Araneidae, 57
Archaea, 659
Archaeognatha, 557
Archaeomerycidae, 222
*Archaeomeryx*, 222
*Archeoglobus fulgidus*, 659
Archosauria, 403, 833
Archosaurs, 239–240, 253–255
*Arctocyon*, 858
*Arctonyx*, 581, 590
  *collaris*, 573, 577, 582
Area of endemism, 271–280, 717–723
Arecaceae, 346
Are the dental data really at odds with the molecular data? Morphological evidence for the whale phylogeny (re)reexamined, Maureen A. O'Leary, John Gatesy, and Michael J. Novacek, 853–866
Argasidae, 300, 309
Argasinae, 300
*Argas lagenoplastis*, 300
*Argia*, 94
  *sedula*, 94
Aris-Brosou, Stéphane, Least and most powerful phylogenetic tests to elucidate the origin of the seed plants in the presence of conflicting signals under misspecified models, 781–793
Arnedo, Miquel, *see* Hormiga, Gustavo, 70–88
*Arthropeas magnum*, 756
Arthropods, 70, 107, 296, 299–300, 304–305, 307, 518, 557
*Artibeus jamaicensis*, 462–464
Articulata, 518, 520
Artiodactyla, 15–22, 724–729, 854–855, 857–858
Artiodactylans, 853, 857, 860
  hippopotamid, 854
  suine, 854
*Ascaltis*, 314–315, 332–333
*Ascandra*, 314–315, 332–333
  *falcata*, 332
  *minchini*, 332
Ascomycetes, 430
Asilidae, 750, 756
Asiloidea, 750, 753
*Asioryctes*, 858
*Aspidelaps*, 451
  *scutatus*, 440
Assessing progress in systematics with continuous jackknife function analysis, Jeremy A. Miller, 55–65
Assumptions, 352–367
Asteraceae, 348, 820, 823
Astigmata, 430
Asymtomatic enumeration, 110–118
*atp*6, 657, 659
*atp*8, 656–659
Atelestidae, 756
*Atelestus pulicarius*, 756
Atherinomorpha, 549
Atractaspididae, 439–440, 455–456
Atractaspininae, 440, 447
*Atractaspis*, 451, 455
  *bibronii*, 440
  *corpulenta*, 440
  *micropholis*, 440
Augochlorini, 23
Australia, 757–780
*Australictus*, 24
*Austrevylaeus*, 24, 26, 30
Autopomorphic characters, 865
Aves, 37, 394, 403–422
Avian DNA, 389
Avila, Luciano J., *see* Morando, Mariana, 159–185
Axial symmetry, 311–333
*Axinella polypoides*, 316
*Azemiops*, 451, 453
  *feae*, 440, 453
Azemiophinae, 440

Badgers, 571–573
  American, 571–573
  Chinese ferret, 573
  Eurasian, 573
  hog, 573
  honey, 572

Old World, 577, 590
stink, 573
Baeriida, 314, 317–318, 320, 323–324, 329–330
Baeriidae, 314, 320, 323
*Balaenoptera*, 209, 211–213, 215
*physalus*, 210
Balaenopteridae, 209–210
*Balaenoptera*, 706, 712
Balanidae, 557
Barker, Nigel P., *see* Kelly, Christopher M., 439–459
Barker, Stephen C., *see* Murrell, Anna, 296–310
Barnacles, 557
Barriel, Véronique, *see* Hugot, Jean–Pierre, 735–744
Basidiomycetes, 525
Basidiomycotina, 430
Bassariscus astutus, 573
Bats, 57, 462, 472
short–tailed, 460
Bayesian, 48 -54
analysis, 26, 30, 89, 96, 102, 133, 140, 152, 165, 167, 181–182, 211, 390, 439–459, 477–487, 571, 576, 583, 604–617, 618, 621, 623, 629, 665, 671, 708, 713–714, 745–756, 760, 797, 801–803, 807
approach, 171, 791
consensus tree, 622
estimation, 131–158, 394, 714, 745
factors, 48–50, 52–54, 649–664
framework, 48–49, 95, 372, 650, 666, 714, 760, 790–791
hypothesis, 100
inference, 48, 386, 442, 444–446, 642, 665–673
information analysis, 377
information criteria (BIC), 675–676, 719
methods, 23, 48, 159, 477, 696–697, 705–716, 726, 748, 801
model, 49, 650, 791
model selection, 674–683
phylogenetics, 48, 54, 89–109, 652
posterior probability, 30, 102, 215, 396, 554, 559, 612–613, 665–673
relaxed clock, 206–228
support, 623
tests, 607, 615
theorem, 48, 136, 267, 650, 655
trees, 30, 100, 167, 455, 457, 589, 750, 762, 764, 767, 771, 773, 802
Bears, 711–713, 794
*Bedfordia*, 91, 94, 96
sp., 102
Bees, 1–14, 23–26
allodapine, 1
corbiculate, 23, 33
halictid, 23–24, 33
honey, 26
Beetles,
carabid, 70, 103
*Ips*, 677
longhorn, 89
Bennettitales, 753
Bennett, Jonathan R., *see* Scotland, Robert W., 539–547
*Bernissartia*, 412
*Beroe*, 317
*cucumis*, 316
Best-fit trees, 4
$\beta$2-microglobulin, 738
$\beta$-casein, 207
$\beta$-tubulin, 596, 598–600
*Bicatillus*, 619, 623, 639
*extinctorum*, 623
Bidirectional sequencing, 2
Bifurcating tree (*see also* binary tree), 642
Bilateria, 311, 316–318, 320
Binary
characters, 244, 247, 256–257, 335, 535
coding, 841
tree (*see also* bifurcating tree), 114
Bininda-Emonds, Olaf R. P., Novel versus unsupported clades: assessing the qualitative support for clades in MRP supertrees, 839–851
Bininda-Emonds, Olaf R. P., Kate E. Jones, Samantha A. Price, Richard Grenyer, Marcel Cardillo, Michael Habib, Andy Purvis, and John L. Gittleman, Supertrees are a necessary not-so-evil: a comment on Gatesy et al., 724–729
Biochemical tree, 403
Biodiversity crisis, 428–435
Biogeographic, 552
analysis, 258, 761, 824
barriers, 618–640
history, 552, 722
inference, 618
pattern, 820
studies, 777
Biogeography, 7, 70–88, 240, 255–258, 430, 509, 618, 622, 717–723, 757–780, 794, 812, 820–841
historical, 271–280
Bioinformatics, 119–124
Biostratigraphy, 240
Biosynthetic pathways, 334–335, 337, 339, 342, 346, 348
Biotic elements, 271–280, 717–723
Biotic element analysis in biogeography, Bernhard Hausdorf and Christian Hennig, 717–723
Birds, 41, 45–46, 230, 283–287, 293, 334, 391, 404, 549–551, 777–778
ratite, 549–553
Birth–death
model, 641, 644–645, 647, 761
process, 132, 641–648
*Bitis*, 451
*arietans*, 440, 459
*Blackburnia*, 103
Black caiman, 388
Black ravens, 353
Black-tailed deer, 210
Blesbok, 210
Blouin, Christian, *see* Susko, Edward, 594–603
Blow flies, 745
*Boa*, 447, 451
*constrictor*, 440, 443, 447
Boaedontinae, 455
Body shape variation, 488–500
Boidae, 440
*Boiga*, 447, 451, 457
*cynodon*, 440
*dedrophila*, 440
Boiginae, 457
Boleichthys, 492–493
Boleostoma, 492–493
Bollback, Jonathan P., *see* Harshman, John, 386–402
Bollback, Jonathan P., *see* Huelsenbeck, John P., 131–158
*Bolyphantes alticeps*, 73, 88
Bombinatoridae, 794
Bombyliidae, 749–750, 754
*Bombylius*, 754
*major*, 756
Bonferroni
adjustment, 723
correction, 16, 279, 686, 783, 807
procedure, 802
*Boophilus*, 299, 305–306, 310
*annulatus*, 299, 306

*decoloratus*, 299
*geigyi*, 299
*kohlsi*, 299
*microplus*, 298–299, 306
Bookstein, Fred, *see* Rohlf, F. James, 66–69
Bootstrap, 3, 39, 41, 43, 95, 99, 128–129, 554, 556, 559, 561, 575–576, 589, 600, 665–673, 737, 791, 841
analysis, 7, 26, 447, 580, 609, 611, 621, 697, 775, 824, 842
consensus, 30, 32, 802
consensus trees, 609
estimates, 599
frequencies, 839, 842, 844, 847
maximum likehood, 667–668
nonparametric, 166, 506, 594, 665–667, 669, 671–672, 726, 760, 790, 801
parametric, 18, 598, 667, 726, 757–780, 781
percentages, 102, 211, 249, 737, 762, 781–793
proportion, 129
RELL, 390
replicates, 42, 44, 477, 479–481, 621, 823
support, 4–8, 26, 30, 42, 55, 62, 95, 167, 218, 444, 446, 467–468, 508, 540, 576–577, 581–583, 608–609, 611, 614, 621, 623, 738, 764, 829, 842, 844
trees, 251, 380–381, 451, 455, 465
value, 5, 7, 45, 95, 101–102, 211, 289, 320, 322, 336, 378, 477, 480, 484, 504, 506, 508, 510, 557, 576, 588, 609–610, 614–615, 665, 667, 727, 738, 748, 760, 802, 826–829
Borchiellini, Carole, *see* Manuel, Michael, 311–333
Boreallodapini, 2
*Borealosuchus* ssp., 409
*Boronia*, 347
*Bos*, 209, 706
*taurus*, 210–212, 405
Boselaphini, 209–210
*Boselaphus*, 209–212
*tragocamelus*, 210
*Bostrycapulus*, 619, 623, 631, 636, 639
cf. *aculeata* sp.1, 631
*Bothriocroton*, 300, 304–307, 309
*concolor*, 300
*undatum*, 300
Bothriocrotoninae, 300, 307
Boulengerina, 447, 451
*annulata*, 440
*fasciatus*, 440
sp., 440
Boury-Esnault, Nicole, *see* Manuel, Michael, 311–333
Bovidae, 206–228
Bovinae, 210, 218–219
Bovini, 209
Brachycera, 745–756
*Brachylinga* sp., 756
Branch-and-bound search, 368, 377, 390
Branchiopoda, 557
Branch and bound search, 479
Branch length, 22, 41, 45, 48–50, 52, 76, 81–82, 96, 127–130, 133–135, 137, 156, 211–212, 223, 358–359, 377, 412, 415, 424, 427, 478, 481, 484–485, 508, 529–531, 535, 575, 589, 596, 608, 621, 629, 641, 644, 651–652, 655, 666–667, 671–672, 675, 692, 696–697, 699–700, 703, 709, 761, 782, 784–785, 791, 820, 825, 827, 831–832, 834–835, 841, 849–851, 865–866
error, 679–681
estimates, 396, 676–677, 682, 747
Branch lengths and support, Mark Wilkinson, François-Joseph Lapointe, and David J. Gower, 127–130
Branch lengths and support: revisited, David A. Morrison, 849–851
Branch support (*see also* bootstrap, clade support, decay, posterior probability), 15, 26, 30, 211, 323, 407, 773
partitioned, 15–22, 76, 79, 407, 409, 412
Branch-swapping, 30, 368–370, 842, 857
algorithm, 287, 371
NNI, 372
TBR, 39, 75, 166, 275, 318, 336, 377–378, 390, 407, 444–445, 463, 504, 506, 575, 607, 621–622, 666, 681, 748, 760, 801, 823–824
*Brassica*, 374
Braun, Michael, *see* Harshman, John, 386–402
*Braunsapis*, 2, 5, 7
*paradoxa*, 2
*protuberans*, 2
*unicolor*, 2
*vitrea*, 2
BRCA1, 605, 613–615
Bremer
analysis, 211
support (*see also* decay), 4, 10–11, 55, 62, 75–77, 127, 307, 318, 320, 322, 554–564, 571, 575, 581–584, 588–589, 665, 842
*Brevineura*, 1, 7–8, 10–11
*ploratula*, 2
*xanthoclypeata*, 2
Brevirostres, 390
Briggs, John C., Fishes and birds: Gondwana life rafts reconsidered, 548–553
Britton, Tom, *see* Erixon, Per, 665–674
Brownian motion parameter, 714
Brown, Rafe M., *see* Evans, Ben J., 794–823
Bryant, David, *see* François-Joseph Lapointe, 865–868
*Bufo valliceps*, 287
*Bungarus*, 451
*fasciatus*, 440
Bull, Nicholas J., *see* Schwarz, Michael P., 1–14
Burn-in trees, 607
Butterflies, 794
nymphalid, 28
swallowtail, 589

Caenogastropod, 620, 623
Caenophidia, 439–459
*Caiman*, 391, 396, 415
*crocodilus*, 405, 412, 417
*latirostris*, 405
*yocare*, 388
Caimanae, 409
*Cairina moschata*,
Calamariinae, 440, 457
Calcarea, 311–333
Calcaronea, 311, 313–315, 318, 320, 322–325, 328–329
Calcinea, 311, 313–315, 318, 320, 322–325, 328–329
Calcispongia, 311–333
Callilictus, 24
*Callimico*, 740, 742
*goeldii*, 736
Calliopheinae, 443
*Callithrix*, 706, 740, 742
*geoffroyi*, 736
*jacchus*, 736, 738
Callitrichinae, 740, 742
*Calloselasma*, 447, 451
*rhodostoma*, 441
*Calyptraea*, 618–640
*aspersa*, 638
cf. *conica*, 638
cf. *lichen*, 638

*chinensis*, 623, 638
*fastigata*, 638
*mamillaris*, 638
Calyptraeid, 620, 629, 631–633
biogeography, 631
evolution, 632
gastropods, 618–619
larvae, 631
taxonomy, 619
Calyptraeidae, 618–619, 623, 632
Camels, 20, 221, 857
Camin–Sokal parsimony, 444, 446
Campbell, Christopher S., *see* Smedmark, Jenny E. E., 374–385
Campbell, Nick, J. H., *see* Murrell, Anna, 296–310
Canary, 389
Canidae, 846
Canids, 707–708
*Canis*, 706
Cannatella, David C., *see* Evans, Ben J., 794–823
Canonical
correspondence analysis (CCA), 687–689, 693
ordination (CO), 687, 693
phylogenetic ordination (CPO), 684, 687–689, 691–694
tree, 113, 115
variate analysis (CVA), 190, 196, 198, 491, 496
Canonical phylogenetic ordination, Norberto P. Giannini, 684–695
*Canthumeryx*, 224
Capreolini, 209
*Capreolus*, 209, 212, 215, 218–219
*capreolus*, 210
Caprini s.l., 209–210, 215, 225
Capulidae, 639
Capulids, 620, 623
*Capulus ungaricus*, 639
Carabid beetles, 70, 103
Cardillo, Marcel, *see* Bininda-Emonds, Olaf R. P., 724–729
Cardini, Andrea, Geometry of the marmot (Rodentia: Sciuridae) mandible: phylogeny and patterns of morphological evolution, 186–205
*Carios capensis*, 300
Carnivore supertree, 845
Carnivora, 571, 839–851
*Carollia perspicillata*, 472
Cartesian transformation grids (CTGs), 488, 491–492, 497
Caryophyllaceae, 820, 832
Cassowaries, 550
*Castor canadensis*, 606
Castoridae, 604–606, 608–611, 613–615
*Casuarius*, 389
*casuarius*, 388
Catarrhini, 738
Catonotus, 492–493
Causinae, 441, 453
*Causus*, 451
*defilippi*, 441
*rhombeatus*, 441
Cattle, 206
CB1, 604, 606–610, 613
Cave salamanders, 501
*Cavia*, 649–664
*porcellus*, 649, 656
Cenozoic, 206, 631, 746
Centipedes, 559
Cephalocarida, 557
Cephalophini, 209–210
*Cephalophus*, 209, 212, 215, 221
*dorsalis*, 210
*maxwelli*, 210
Cerataphidini, 137, 139
*Ceratina*, 8
*japonica*, 2, 5
Ceratinini, 1, 2, 9
*Cercomegistus* sp., 300
*Cercopithecus*
*hamlyni*, 736
*nictitans*, 736
*Cerberus*, 451
*rhynchops*, 440, 446
*Cermatobius*, 559
Cervidae, 206–207, 209–210, 212–213, 215, 217–222, 224
Cervinae, 209
*Cervus*, 207, 209, 212, 215, 218–219
*elaphus*, 210
*unicolor*, 210
Cetacea, 15–22, 209, 221, 724–729, 853–855, 857–858, 860
Cetaceans, 853, 860
phylogeny, 854
Cetartiodactyla, 206, 221
*Chaetodipus*, 609, 614
*californicus*, 606
*Chalinolobus tuberculatus*, 462
Characters, 239–252, 851–852, 864
adult, 8–9
anatomical, 742
autopomorphic, 865
binary, 244, 247, 256–257, 335, 535
chromatic, 178
cladistic, 854
coding, 334–351, 401, 405, 540
complex, 245
constant, 378
construction, 239–241, 243, 246, 253
convergent, 502, 510
correlation, 131–158
cranial, 187
cytological, 311, 313, 326–328
dental, 738, 853, 857, 860
diagnostic, 323, 328, 545
discrete, 136, 825
distribution, 264–265
DNA, 56, 348, 405, 509
embryological, 311, 313, 327–328
equally weighted, 318, 327, 358
evolution, 56, 152, 263, 318, 328, 364, 416, 543, 647, 833
foot, 858
fruit, 374–375
gap, 317–318
genitalic, 8, 91
homoplastic, 589, 767
indel, 389, 823, 826, 829
informative, 16, 378, 575, 583
larval, 8–9, 92
life-history, 9
mandibular, 192
mapping, 27, 131–158
meristic, 506, 513
molecular, 55, 70, 76–77, 131, 199, 209, 296, 298, 409, 738, 742
morphological, 55, 57, 70, 72, 74, 87–88, 89, 91, 131–132, 135, 142, 178, 181, 199, 209, 213, 217, 219, 298, 313–314, 317, 323, 327–328, 340, 389–390, 400, 406–407, 409, 412, 420, 439, 442, 457, 474, 501–503, 506, 509–510, 537, 539–543, 545, 559, 572, 604, 632, 738, 742, 748, 760, 855–857

multistate, 26, 75, 239, 255–257, 335
osteological, 187, 760
parisitological, 738, 742
parsimony-informative, 4, 298, 323, 345, 446, 451, 529, 531, 773
phylogenetically informative, 55–56, 826
polymorphic, 505, 833
primary, 725–726
pupal, 8
reductive, 245
reweighted, 318, 327, 371
shell, 619, 623
skeletal, 197, 199, 222, 313, 403
snout, 416, 621–622
state, 87, 132, 149, 239, 245–246, 250, 253, 296, 309–310, 352, 354, 357, 400, 405, 427, 502–503, 505, 542, 760, 831, 835
summary, 337, 342
systematic, 243
two-state, 236
unambiguous, 539, 557
uninformative, 76, 217, 575, 581, 826
vertebral, 439
*Charadrius semipalmatus*, 286
*Cheilea*, 632, 639
*equestris*, 639
Cheirogaleidae, 711–713
*Cheirogaleus*, 706
Chelicerata, 56, 296, 557
Chevrotains, 206
lesser Malay, 210
Chicken, 388–389, 391, 551
*Chilalictus*, 24
Chilopoda, 557
*Chimeromyia*, 749
Chimpanzee, 52–53, 711–713, 715
*Chinchilla laniger*, 606
Chinchillidae, 606
Chinese
muntjak, 210
water deer, 210, 220
Chipmunks, 194, 197
Chippindale, Paul T., *see* Wiens, John J., 501–514
Chirononomidae, 430
Chi-square tests, 307, 395, 397–398, 607
Chloroplast
DNA, 374–375, 382
genome, 336
phylogeny, 375
Choanoflagellata, 316–317
*Chordeiles acutipennis*, 286
Chromatic characters, 178
Chromatograms, 94
*Chrysops carbonarius*, 756
Cichlidae, 549, 552
Cichlid fish, 549–553
*Ciconia nigra*, 286
*Cirripathes*, 317
*lutkeni*, 316
Cirripedia, 557
Clades, 11, 15–17, 19–20, 23, 34, 55, 62, 70, 76–77, 79, 81, 100–101, 103, 105–106, 119, 122, 125, 140, 159, 165–167, 172, 178–179, 186–187, 220, 241, 249, 275, 278, 304–305, 318, 320, 323–324, 327, 329, 334, 344, 374–375, 377, 379, 383–384, 390, 403, 409, 439, 445–447, 452, 455, 463, 468, 494, 503, 508, 511, 529–531, 535, 542, 557, 560–561, 581, 583, 585, 590, 604, 614, 618, 623, 629, 631–632, 665, 667–669, 671–672, 688, 697, 711–712, 724, 749, 761–762, 767, 771, 776, 797, 802, 804, 807, 810–811, 827, 829, 832, 839–840, 843–847
phylogenetic, 714
Cladistic
analysis, 8, 81, 253, 403, 541, 572, 738
characters, 854
information content (CIC), 241
hypothesis, 92
relationship, 70–71
parsimony, 356, 362, 365
Cladistics, 63, 70–88, 520
molecular, 187
Cladogenesis, 641–642, 644–645, 647, 746, 757, 761, 834
Cladogenetic model, 641
Cladogram, 5, 70, 73, 76–77, 79, 81, 83–84, 195, 255–258, 272, 359, 362, 406–407, 520, 525–526, 548–549, 684, 722, 742, 802
coding, 255–259
molecular, 187, 742
*Claravis pretiosa*, 38
Clarifying Aetosaur phylogeny requires more fossils, not more trees—reply to Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), Andrew B. Heckert and Spencer G. Lucas, 253–254
*Clarkia*, 347
*Clathrina*, 314–315, 320, 329, 332–333
*cerebrum*, 316
*clathrus*, 329
Clathrinida, 314, 318
Clathrinidae, 314
Clayton, Dale H., *see* Johnson, Kevin P., 37–47
*Climacoceras*, 224
Clocklike trees, 233
Clock model (*see* molecular clock)
Cluster analysis, 194, 205, 497, 722
*C-mos*, 758, 762, 764
*C-myc*, 386–402
Cnidaria, 311, 316–317, 320, 430, 520
*Coahomasuchus*, 241, 243, 250–251, 254
Coding, 239–252, 851–852
cladograms, 255–259
indels, 444, 575
Codons/codon positions, 3, 5, 7, 10, 20, 28, 391
Cody, Jeannine H., *see* Rodman, James E., 428–435
Coelenterates, 518
Coenagrionidae, 89, 91–92
*Coenomyia ferruginea*, 756
Coevolution, 37–47
Coleoptera, 430
Collin, Rachel, Phylogenetic relationships among Calyptraeid gastropods and their implications for the biogeography of marine speciation, 618–640
Colonization, 70–88
*Coluber*, 451
*constrictor*, 440
Colubridae, 440, 456
Colubrid snakes, 758
Colubrinae, 439–440, 447, 452, 455, 457
Colubrini, 457
Colubroidea, 439–459
*Columba*
*guinea*, 38
*livia*, 38, 286
*Columbicola*, 37–39, 41–42, 45
*adamsi*, 38
*bacillus*, 38
*baculoides*, 38
*columbae* 1, 38

*columbae* 2, 38, 42
*exilicornis*, 38
*extinctus*, 38, 42
*gracilicapitis*, 38, 42
*gymnopeliae*, 38
*macrourae* 1, 38, 46
*macrourae* 4, 38
*macrourae* 5, 38
n. sp., 38
*passerinae* 1, 38, 42
*passerinae* 2, 38
*theresae*, 38, 46
*timmermanni*, 38
*veigasimoni*, 38
*xavieri*, 38
Columbiformes, 37–47
*Columbina*, 46
*inca*, 38, 46
*passerina*, 38, 46
*Coluria*, 374, 381–383
*geoides*, 374, 376, 381–382
Colurieae, 374, 377–378, 382
Combinatorics of tandem duplication trees, Olivier Gascuel, Michael D. Hendy, Alain Jean-Marie, and Robert McLachlan, 110–118
Combined data, 55–65
analysis, 96
evidence, 403–422
tree, 612
Combined support for wholesale taxic atavism in gavialine Crocodylians, John Gatesy, George Amato, Mark Norell, Rob DeSalle, and Cheryl Hayashi, 403–422
Combining data, 528–538
Comparative
biology, 55–65
method, 23–36
Comparing bootstrap and posterior probability values in the four–taxon case, Michael P. Cummings, Scott A. Handley, Daniel S. Myers, David L. Reed, Antonis Rokas, and Katarina Winka, 477–487
Comparison of likelihood and Bayesian methods for estimating divergence times using multiple gene loci and calibration points, with application to a radiation of cute-looking mouse lemur species, Ziheng Yang and Anne D. Yoder, 705–716
Complex characters, 245
Component coding, three-item coding, and consensus methods, David M. Williams and Christopher J. Humphries, 255–258
*Compsomelissa*, 1–2, 4–5, 7–12
*borneri*, 2
Computer simulation (*see also* simulation and Monte Carlo simulation), 528–538, 696–704
Computing the uniform component of shape variation, F. James Rohlf and Fred Bookstein, 66–69
Conflict, 839–851
Congruence, 37, 55–65, 193, 329, 539–548, 555, 607, 613, 669, 717, 865–868
Conifers, 784
Consensus, 255–259, 849–851
efficiency (CE), 241
trees, 55, 128, 140, 241, 251, 255, 258, 344, 450, 531, 850, 855, 865–868
Consistency index, 3, 5–6, 10, 28, 76, 240, 247–248, 252, 304, 339, 345–346, 348, 392, 396, 577, 581–583, 762, 764, 826
Constrained tree, 16–18, 20, 22, 407, 761, 807, 847
Continuous
jackknife function analysis (CJF), 55–58, 61–63
time Markov model, 131–134, 147, 149, 651
time Markov process, 179, 644
Control region (CR), 460–476
*Conus*, 619
Convergence, 501–514, 614
sciurid, 185–205
Convergent
characters, 502, 510
evolution, 296, 497
Conway, Lindsay, *see* Danforth, Bryan N., 23–36
Co-occurrence, 271–280
Cooper, Steven J. B., *see* Schwarz, Michael P., 1–14
Copath
analysis, 41
tree, 45
Cophylogenetic analysis, 737
Cophylogeny, 735–744
*Coracias caudata*, 286
*Coragyps atratus*, 388
Corbiculate bees, 23, 33
Correspondence analysis, 687
Corroboration, 352–367
*Cosmiomma*, 305–306
Cospeciation, 37–47
*Coturnix pectoralis*, 286
Cows, 210
Crangonyctidae, 430
Cranial characters, 187
Credibility regions (CR), 390, 394
*Crepidula*, 618–640
*aculeata*, 636
*adunca*, 637
aff. *depressa*, 636
aff. *marginalis*, 637
aff. *onyx*, 638
aff. *williamsi*, 637
*arenata*, 637
*argentina*, 636
*atrasolea*, 636
*cerithicola*, 637
cf. *aculeata*, 636
cf. *aculeata* Cape Verde, 623, 636
cf. *aplysioides*, 637
cf. *incurva*, 637
cf. *nivea*, 637
cf. *onyx*, 638
cf. *perforans*, 637
*complenata*, 629, 631, 637
*convexa*, 629, 637
*coquimbensis*, 636
*costata*, 636
dd. *onyx*, 638
*depressa*, 636
*excavata* Mexico, 620, 632, 637
*excavata* Peru, 620, 637
*fimbriata*, 636
*fornicata*, 619, 632, 637
*gibbosa*, 637
*grandis*, 632, 637
*gravispinosa*, 636
*immersa*, 636
*incurva*, 632, 637
*lessoni*, 636–637
*maculosa*, 638
*marginalis*, 638
*monoxyla*, 636
*naticarum*, 638
*navicula*, 638

*norrisarum*, 638
n. sp., 638
n. sp. dd., 638
n. sp. pt., 638
*nummaria*, 637
*onyx*, 619, 631, 638
*perforans*, 637
*philippiana*, 623, 637
*plana*, 637
*porcellana*, 629, 638
*protea*, 637
s.l. , 637
s.s., 623, 631–632, 636
*striolata*, 637
*unguiformis*, 637
*ustulatulina*, 638
*williamsi*, 623, 637
*Crepipatella*, 623, 631, 636
*capensis*, 631–632, 636
*dilatata*, 632, 636
*fecunda*, 632, 636
*lingulata*, 636
n. sp. , 636
Cretaceous, 23, 33, 403, 419, 550–552, 631, 746, 753, 861
early-, 548
mid-, 549, 552
late-, 419, 549–552, 753
upper-, 753
*Cretaformia*, 749
Crickets, 70, 833
Crocodile, 285
African slender-snouted, 388
Crocodilians, 240, 283, 286, 293, 389, 403
Crocodylia, 386–402, 403–422
Crocodyliformes, 386, 415
Crocodylidae, 386, 392, 394
Crocodylinae, 403, 405, 408–409, 412, 415, 417, 419
Crocodyloidea, 391
*Crocodylus*, 386, 391, 396
*cataphractus*, 388, 391, 405, 408–409, 415, 417
*intermedius*, 405, 408, 417
New World, 409
*niloticus*, 285
*rhombifer*, 405, 408
Crotalinae, 441, 447, 453
*Crotalus*, 447, 451
*horridus*, 441
*viridis*, 441
*Crotaphopeltis*, 447, 451, 457
*hotamboeia*, 440
*Crucibulum*, 618–640
*auricula*, 638
cf. *personatum*, 638
cf. *serrata*, 638
*concamaratum*, 623, 638
*lignarum*, 638
*radiata*, 638
*scutellatum*, 638
*spinosum*, 638
*tenuis*, 638
*Crunomys*, 812
Crustacea, 430, 432, 557
*Crystallaria*, 496–498
*asprella*, 491, 496–499
Ctenodactylidae, 605–606
Ctenohystric, 605, 615
*Ctenonomia*, 24
Ctenophora, 311, 316–318, 320, 327–328
*Cuculus pallidus*, 286
*Culex pipiens*, 745
Cumaceans, 430
Cummings, Michael P., Scott A. Handley, Daniel S. Myers, David L. Reed, Antonis Rokas, and Katarina Winka, Comparing bootstrap and posterior probability values in the four-taxon case, 477–487
Culicomorpha, 754
CXCR4/CCR5 coreceptor, 649–664
Cycads, 784
*Cyanoramphus*, 471
*Cyclorrhapha*, 746, 749–751, 753
*Cymatium parthenopeum*, 640
*Cynomys*, 189, 194
*leucurus*, 204
*ludovicianus*, 189, 204
*Cypraea spurca*, 640
Cyprinodontiformes, 548–459
*Cypripedium*, 346
Cytogenetic analysis, 374
*Cytochrome b*, 2–5, 7–8, 10–11, 16–18, 20, 22, 39, 41–42, 161, 163–166, 171, 179, 181, 186, 193, 195, 198–199, 206–207, 209–212, 215, 297, 443, 445–447, 451, 455, 471–472, 488, 504, 508, 571–593, 605, 657, 659, 677, 680–681, 705–706, 709, 727, 861
Cytochrome oxidase I (COI), 1–5, 7–8, 10–11, 23, 26–30, 32, 37, 39, 41–42, 45, 72–73, 139, 297–298, 557, 559, 618–640, 657–659, 674
Cytochrome P-450, 206, 209–212, 219
Cytochrome oxidase II, 33, 94, 95, 139, 207, 705–706, 709
Cytological characters, 311, 313, 326–328

*Dactylanthus taylorii*, 473
*Dalloapula*, 9
*Damaliscus*, 209, 212, 215, 218
*pygargus*, 210
Damselflies, 70, 833
*Megalagrion*, 89–109, 833
Danforth, Bryan N., Lindsay Conway, and Shuqing Ji, Phylogeny of eusocial *Lasioglossum* reveals multiple losses of eusociality within a primitively eusocial clade of bees (Hymenoptera: Halictidae), 23–36
Darters, 488–489, 494, 498
*Dasypeltis*, 447, 451, 457
*medici*, 440
*scabra*, 440, 459
Data
combination, 439–459
conflict, 386–402
non-independence, 724–729
partitions, 386–402
visualization, 55–65
*Daubentonia*, 706, 712
Dayrat, Benoît, Roots of phylogeny: how did Haeckel build his trees?, 515–527
DeBry, Ronald W., Identifying conflicting signal in a multigene analysis reveals a highly resolved tree: the phylogeny of Rodentia (Mammalia), 604–617
Decay
analysis (*see also* Bremer support), 12, 128, 243, 407
index, 128, 336, 377, 381, 842
Decision theory, 674–683
Deduction, 259–271
Deep sea mollusks, 430
Deer,
black-tailed, 210

Chinese water, 210
European roe, 210
mouse, 206, 854, 857
musk, 206–207, 210, 219–220
red, 210
Dei-Cas, Eduardo, *see* Hugot, Jean-Pierre, 735–744
Deletion (*see* indel)
*Delma*, 757–758, 762, 764, 767, 770–771, 773, 775, 777–778
(*Aclys*) *concinna*, 759, 762, 764, 770, 775, 777–778
*australis*, 759, 775, 778
*borea*, 759, 775, 778
*borea haroldi*, 759
*butleri*, 759, 770, 778
*elegans*, 758, 775
*fraseri*, 757–759, 762, 775–778
*fraseri fraseri*, 762, 775–776
*fraseri petersoni*, 762, 775–776
*grayii*, 757, 759, 762, 775–778
*impar*, 759, 775, 778
*inornata*, 759
*labialis*, 759, 764, 770, 775, 777–778
*mitella*, 759, 762, 770
*molleri*, 759, 775, 778
*nasuta*, 759, 778
*pax*, 759, 762, 775
*petersoni*, 757, 759, 777
*plebeia*, 758, 775
*tincta*, 759, 775, 778
*torquata*, 759, 775, 778
Demanche, Christine, *see* Hugot, Jean-Pierre, 735–744
Demospongiae, 311, 314, 316–317, 320, 328–330
Dendrelaphis, 447, 451, 457
*calligastra*, 440
*pictus*, 440
*Dendroaspis*, 447, 451, 456
*polylepis*, 441
sp., 441
Dendrogram, 192–193, 195, 199
mandibular, 195, 197
UPGMA, 193
Dendroica, 776
Dental characters, 738, 853, 857, 860
Dentition, 16
*Dermacentor*, 299, 305–306, 310
*andersoni*, 299
*marginatus*, 299
*reticulatus*, 299
s.l. , 299, 305
*variabilis*, 299
DeSalle, Rob, *see* Gatesy, John, 403–422
*Desmatosuchus*, 241–243, 248–252, 254
*haplocerus*, 254
de Queiroz, Kevin and Steven Poe, Failed refutations: further comments on parsimony and likelihood methods and their relationship to Popper's degree of corroboration, 352–367
DHPS, 735–738, 740, 742
*Diacodexis*, 858
Diagnostic characters, 323, 328, 545
Dialictus, 24, 28, 32–33
*Diaphanoeca grandis*, 316
Diapsida, 239, 253
Dicotyledons, 524
*Dicrocerus*, 224
*Didelphis*, 706, 708
*virginiana*, 285, 709
Diesmos, Arvin, *see* Evans, Ben J., 794–823
Dinosauria, 839
Dinosaurs, 253
*Diospyros sandwichensis*, 820
*Dipodomys*, 614
*spectabilis*, 606
*Dipsas*, 447, 451
*Dinodon*, 447, 451
*semicarinatum*, 440, 445
*Diogmites* sp., 756
Diphyllidae, 430
Diplodactyline geckos, 758, 760
*Diplodactylus*, 757
*damaeus*, 758–759
*tessellatus*, 758–759
Dipodidae, 604–606, 608, 613–614
*Dipsas catesbyi*, 441
Diptera, 430, 557, 745–756
*Dipus*, 614
*sagitta*, 606
Dirichlet
density, 711
distribution, 607, 652, 655, 749
Discete characters, 136, 825
Discrete-gamma
distribution (*see also* gamma distribution), 165, 445, 666, 698, 700, 703
estimate (DGE), 595–596, 599–601
probability estimate (DGPE), 596–602
Dispersal, 549–553, 794–819
Distance analysis, 334, 343
Divergence time, 705–716, 745–756
DNA, 2–3, 74, 94, 110, 166, 212, 221, 298, 306, 334, 340, 342, 345, 370, 382, 389, 391, 408–409, 419, 506, 509, 551–552, 606, 620, 725, 727, 760, 857
analysis, 551
avian, 389
characters, 56, 348, 405, 509
chloroplast, 374–375, 382
evolution, 443
extraction, 26, 38, 462, 736
fingerprints, 404
genomic, 161, 315–316, 443, 550, 566, 574, 606, 758, 823
hybridization, 550–551, 573
isolation, 375
mitochondrial, 3–4, 7, 12, 37, 48, 52, 89, 94–101, 103, 159–185, 296, 386–387, 395, 407–408, 415, 418, 423, 439–459, 460–461, 464, 471, 474, 503–504, 506, 508–509, 548, 551, 571–572, 583, 586, 589, 656, 674, 677, 680–681, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776, 794, 798, 801, 808, 810–812
nuclear, 37, 89, 95, 99, 103, 221, 374–375, 382–383, 407–408, 415, 418, 423, 509, 571–593, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776
plasmid, 316
*Pneumocystis*, 735
polymerase, 606
sequence, 38–39, 49, 79, 133, 135, 140, 159, 209, 296, 298, 304, 334–336, 339–340, 344, 346, 348, 387, 396, 400, 404–405, 418–419, 477, 479, 488, 571, 574, 604, 619, 621, 623, 632, 651, 666, 735, 823, 857
sequence data, 23, 509, 530–531, 539, 542, 545, 674, 708, 758, 773, 849
substitution, 135, 140, 575, 644, 647, 824
substitution model, 444
tree, 770, 773, 775
Dog, 591, 711–713
Dolichopodidae, 756
*Dollosuchus*, 416

Dolphins, 501
Donnellan, Stephen, *see* Jennings, W. Bryan, 757–780
Douzery, Emmanuel J. P, *see* Hassanin, Alexandre, 206–228
Doves, 37, 41, 45
  New World ground, 39
*Draco*, 810
*Dremotherium*, 224
*Dromaius novaehollandiae*, 286
*Drosophila*, 2, 89, 383, 557, 745–746, 751, 753–754, 833
  *melanogaster*, 94, 745, 756
  *yakuba*, 94, 298
Drosophilidae, 756
Drosophilid flies, 70
Dryadeae, 374
*Dryomys nitedula*, 606
Dwarf caiman, 388

Echinodermata, 296, 520
*Echinometra* sea urchins, 631
Echmoidea, 288
*Echthralictus*, 24
Ecology, 430, 515, 633, 684
Ecomorphology, 498, 684–695
EF-1$\alpha$, 2–4, 8, 10–11, 23, 26–30, 32, 39, 41–42, 94–99, 101–102, 557, 596, 598, 677, 680
Effect of phylogeny on interspecific body shape variation in darters (Pisces: Percidae), J. Michael Guill, David C. Heins, and Craig S. Hood, 488–500
$\varepsilon$-globulin, 738
*Eichhornia*,
  *azurea*, 154, 156
  *crassipes*, 154, 156
  *diversifolia*, 154, 156
  *heterosperma*, 154, 156
  *meyeri*, 154, 156
  *paniculata*, 154, 156
  *paradoxa*, 154, 156
  sp., 154, 156
Eigenvectors, 69, 694
Eigenvalues, 67
*Eira*, 577, 581, 590
  *barbara*, 573, 581
*Elaphe*, 451
  *flavolineata*, 440
  *obsoleta*, 440
Elapidae, 439, 441, 443, 447, 455–456
Elapinae, 441, 447
*Elapsoidea*, 451
  *nigra*, 441
Elephants, 794
  African, 591
*Ellipsoptera*, 776
Embriological characters, 311, 313, 327–328
Empididae, 754
Empidoidea, 746, 751
*Empis*, 754
Emu, 550–551
*Enallagma*, 94
  *geminatum*, 94
Endemism, area of, 271–280, 717–723
*Enhydra*, 590
  *lutris*, 573, 583, 590
*Enhydris*, 446, 451
  *enhydris*, 440
  *plumbea*, 440
Eocene, 222, 224, 403, 419, 857–858
  early-, 795
  late-, 222, 224
  mid-, 218, 222, 224, 550
*Eoconodon*, 858
*Eogavialis*, 401
*Eotragus*, 225
Ephemeroptera, 106
*Ephydatia muelleri*, 316
Equal-rates Markov (ERM), 229–238
  model, 57
  trees, 233
Equally
  parsimonious tree, 528, 530
  weighted character, 318, 327, 358
  weighted parsimony, 620–621
  weighted tree, 211
*Equus*, 706, 857
Eremoneura, 746, 750–751, 754
*Erethizon dorsatum*, 606
Erethizontidae, 606
Eriksson, Torsten, *see* Smedmark, Jenny E. E., 374–385
Erixon, Per, Bodil Svennblad, Tom Britton, and Bengt Oxelman, Reliability of Bayesian posterior possibilities and bootstrap frequencies in phylogenetics, 665–673
Ermine, 573
*Erythrocoma*, 374
  *triflora*, 374, 376, 379, 383
*Esastigmatobius*, 559
*Escherichia*, 660
  *coli*, 209, 383, 659
Estimation of rates-across-sites distributions in phylogenetic substitution models, Edward Susko, Chris Field, Christian Blouin, and Andrew J. Roger, 594–603
*Etheostoma*
  *blennioides*, 497, 500
  *blennius*, 491, 500
  *caeruleum*, 498, 500
  *coosae*, 492, 500
  *crossopterum*, 491, 500
  *exile*, 498, 500
  *flabellare*, 500
  *fusiforme*, 498, 500
  *inscriptum*, 497, 500
  *jordani*, 492, 498, 500
  *juliae*, 491–492, 498
  *lynceum*, 494, 497, 500
  *nigrum*, 494, 497–498, 500
  *olmstedi*, 491–492, 494, 500
  *proeliare*, 491, 500
  *punctulatum*, 500
  *radiosum*, 493–494
  *simoterum atripinne*, 492, 494
  *simoterum simoterum*, 492, 494
  s.l., 491, 497, 500
  *spectabile*, 500
  s.s., 492–493, 500
  *stigmaeum*, 497–498, 500
  *whipplei*, 491, 493–494, 498, 500
  *zonale*, 491, 494, 500
Etheostomatinae, 488
Eubacteria, 659
Eucalyptus trees, 777
Euclidean
  distances, 67, 337, 676
  geometry, 67
  space, 66–67

*Euconolylops*, 8–9
*Eugeum*, 383
*Eulemur*, 706
  *macaco*, 736
*Eulonchus* sp., 756
Eumeryx, 224
Eumetazoa, 311, 318, 327, 330
Eumycota, 316–317
*Eurycea*, 501–514
  *bislineata*, 505
  *chisholmensis*, 505, 513
  *latitans*, 504–506, 509, 513
  *longicauda*, 505
  *multiplicata*, 504–505, 513
  *nana*, 505, 509, 513
  *naufragia*, 504–506, 513
  *neotenes*, 505, 509, 513
  *pterophila*, 504–505, 509, 513
  *quadridigitata*, 505
  *rathbuni*, 501, 504–506, 508–510, 513–514
  *robusta*, 504, 509
  *sosorum*, 505, 509, 513
  sp., 509, 513
  *tonkawae*, 504–505, 513
  *tridentifera*, 501, 504–506, 508–510, 513–514
  *troglodytes*, 504–505, 508–509, 513
  *wallacei*, 505, 509,
  *waterlooensis*, 504, 509
  *wilderae*, 505
European roe deer, 210
Eusocial evolution, 23
Eusociality, 12–13, 23–36
*Eutamias*, 205
  *sibiricus*, 189, 205
Eutherian mammals, 249, 585, 860
*Euthocodon arambourgii*, 417
Euthyneura, 328
Evans, Ben J., Rafe M. Brown, Jimmy A. McGuire, Jatna Supriatna, Noviar Andayani, Arvin Diesmos, Djoko Iskandar, Don J. Melnick, and David C. Cannatella, Phylogenetics of fanged frogs: testing biogeographical hypotheses at the interface of the Asian and Australian faunal zones, 794–823
Evans, Rodger C., *see* Smedmark, Jenny E. E., 374–385
Even-toed ungulates, 724
Evolution, 1–2, 12, 20, 28, 62, 89, 92, 101–103, 126, 149, 165, 233, 239–252, 254, 296–310, 311–333, 349, 368, 374, 377, 383, 451, 511, 515–527, 549, 551, 573, 576, 645, 658, 660, 725, 735, 740, 851–852
  calyptraeid, 632
  character, 152, 263, 318, 328, 364, 416, 543, 647, 833
  convergent, 296, 497
  DNA, 443
  eusocial, 23
  HIV, 649–664
  island, 820–841
  mammalian, 206
  minimum, 121, 287, 478, 489, 579–580
  molecular, 38, 41, 591, 594–603
  morphological, 186–205, 404, 542
  parallel, 296, 307
  protein, 602
  punctuational, 831
  rate of, 602, 641, 646–647, 677, 746, 833
  reticulate, 374–385
  social, 1–14, 23–36
Evolution of the strategy of long-branch subdivision to improve the accuracy of phylogenetic methods, Steven Poe, 423–427
Evolutionary
  biology, 501–502, 515, 565
  branches, 849
  characters, 404
  character transformation, 354
  convergence, 853
  divergence, 119, 572
  ethnology, 684
  history, 197, 328, 417, 423, 474, 518, 526, 861
  hypotheses, 159
  model, 126, 352, 364, 377, 615, 665, 668, 672, 692, 696–697, 705, 727, 781–782, 788, 791–792, 801, 829, 850
  pattern, 852
  plasticity, 329
  processes, 89, 211, 861
  radiations, 70, 89, 207
  rate, 218, 287, 396–397, 401, 700, 705–716, 748, 753
  studies, 727
  theory, 264–265, 515, 524–525
  trees, 526, 602
Evylaeus, 24, 32–33
*Exoneura*, 1, 12
  *angophorae*, 2
  *robusta*, 2
*Exoneurella*, 1, 7–8, 10–11, 32
  *eremophila*, 32
  *setosa*, 2, 32
  *tridentata*, 2
Exoneuridia, 2, 10
Exoneurines, 1, 9–10
Exons, 376, 383, 391–392, 394
External transcriber spacer (ETS), 820–841

Failed refutations: further comments on parsimony and likelihood methods and their relationship to Popper's degree of corroboration, Kevin de Queiroz and Steven Poe, 352–367
*Fallugia*, 374
False discovery rate (FDR), 783, 790
Falsification, 259–271
Fanged frogs, 794
*Farancia*, 447, 451, 455
  *abacura*, 441
*Farrea occa*, 316
*Fejervarya*
  *limnocharis*, 799, 815
  *vittigera*, 799
Felidae, 845–846
Felids, 707–708
*Felis*, 706
Felsenstein
  model, 463, 703, 708–709, 711–715, 748, 850
  zone, 229, 426, 641–648, 675
FES, 574–577, 581, 583
FIB7, 39, 41–42
*Ficus carica*, 349
Field, Chris, *see* Susko, Edward, 594–603
Filicinae, 524
Fin whale, 210
Fishes, 286, 491, 524, 619
  aplocheiloid, 548–553
  cichlid, 549–553
  cyprinodontiform, 549
  euryhaline, 548
  freshwater, 548–549
Fisher, 573

Fishes and birds: Gondwana life rafts reconsidered, John C. Briggs, 548–553
Fitch
  algorithm, 234
  parsimony, 444, 446
Fitch–Margoliash methods, 290–291, 293
Flatworms, 307
Flies,
  blow, 745
  drosophilid, 70
  flower, 745
  fruit, 557, 745
  hippoiboscid, 37
  horse, 745–746
  house, 745
  snipe, 745–746
  soldier, 746
  stalk-eyed, 32
  xylophagid, 745
Flower flies, 745
Flowers, 336, 341, 343, 346–347, 349, 753
Flowering plants, 820
Foot characters, 858
Fossils, 23, 48, 61, 253–255, 264, 377, 382, 403–422, 521, 524, 528–538, 614, 631, 707, 711, 713, 746–747, 751, 761, 835, 853, 855
  amber, 746
Fossombroniineae, 430
*Fossorella* sp., 639
Four o'clocks, 334–351
Fractional weighting, 258
Fragrance, 334–351
Frequent inconsistency of parsimony under a simple model of cladogenesis, John P. Huelsenbeck and Katherine M. Lander, 641–648
Frogs, 287, 334, 794
  fanged, 794
Fruit, 383
  characters, 374–375
  flies, 557, 745
*F*-test, 687
Fungi, 317, 429–430
*F*-values, 692

G6PD, 738, 742
*Galago*, 706
*Galbula pastazae*, 286
*Galictis*, 590
  *vittata*, 573–574, 576, 581, 589–590
Galliformes, 551
*Gallirex porphyreolophus*, 286
*Gallus gallus*, 286, 396, 405, 408
Gamma
  correlation, 727
  distribution (*see also* discrete-gamma distribution), 4, 42, 50, 95, 135–136, 293, 318, 377, 390, 463, 506, 576, 594–595, 601, 606, 608, 702, 709, 711, 748, 750, 801, 823
  model, 594–603, 750
  shape parameter, 621, 711
$\gamma$-fibrogen, 207
*Gammarus minus*, 509
Gamopetalae, 524
Gap character, 317–318
Gascuel, Olivier, Michael D. Hendy, Alain Jean-Marie, and Robert McLachlan, Combinatorics of tandem duplication trees, 110–118
Gascuel, Olivier, *see* Guindon, Stéphane, 696–704
Gastraea, 311
Gastraeades, 520
Gastropoda, 328
Gastropods, 619, 631
  Calyptraeid, 618–619
*Gastropyxis*, 447, 451
  *smaragdina*, 440
Gatesy, John, George Amato, Mark Norell, Rob DeSalle, and Cheryl Hayashi, Combined support for wholesale taxic atavism in gavialine Crocodylians, 403–422
Gatesy, John, *see* O'Leary, Maureen A., 853–866
Gavial, 403–422
Gavialidae, 398–399
Gavialinae, 403, 416–418, 420
*Gavialis*, 386, 388, 391–392, 394, 396–400, 403
  *gangeticus*, 386, 388, 390, 398, 403, 405–406, 408–409, 412, 415–419
Gavialoidea, 401
*Gavialosuchus*
  *americanus*, 412
  *eggenbergensis*, 417
*Gazella*, 209, 212, 215
  *granti*, 210, 221
  *thomsoni*, 210
Gazelle, 210
Geckos, 758
  diplodactyline, 758, 760
Geinae, 374–379, 382–384
Gekkonidae, 757
Gelocidae, 206
Genealogical trees, 515, 517, 521, 524, 526
Generalized least squares, 491
General-time-reversible (GTR) model, 3–4, 7–8, 18–19, 26, 28–30, 42, 95, 97, 165–166, 212, 286, 318, 390, 392, 396, 445, 463, 576, 579, 585, 608–611, 613, 666–669, 674, 677, 679, 681–682, 737, 762, 784, 801–802, 825, 831, 834
Genes, 296, 605, 609, 612, 652, 658
  A2AB, 605, 614–615
  A6, 674
  *accD*, 336
  $\alpha$-lactalbumin, 206, 209–212, 215, 218–220
  APOB, 574–577, 581, 583
  *atp6*, 657, 659
  *atp8*, 656–659
  $\beta$2-microglobulin, 738
  $\beta$-casein, 207
  $\beta$-tubulin, 596, 598–600
  BRCA1, 605, 613–615
  CB1, 604, 606–619, 613
  CHRNA1, 574–577, 581, 583
  cMDH, 596, 598
  *c-mos*, 758, 762, 764
  *c-myc*, 386–402
  COI, 2–5, 7–8, 10–11, 23, 26–30, 32, 37, 39, 41–42, 45, 72–73, 139, 297–298, 557, 559, 618–640, 674
  COI-1, 1, 5, 657–659
  COI-2, 5, 657–659
  COI-3, 5, 657–659
  COII, 33, 94, 95, 139, 207
  *Cytochrome b*, 2–5, 7–8, 10–11, 16–18, 20, 22, 39, 41–42, 161, 163–166, 171, 179, 181, 186, 193, 195, 198–199, 206–207, 209–212, 215, 297, 443, 445–447, 451, 455, 471–472, 488, 504, 508, 571–593, 605, 657, 659, 677, 680–681, 705–706, 709, 727
  cytochrome P-450, 206, 209–212, 215, 219
  DHPS, 735–738, 740, 742
  EF1$\alpha$, 2–4, 8, 10–11, 23, 26–30, 32, 39, 41–42, 94–99, 101–102, 557, 596, 598, 677, 680

ε-globulin, 738
eRF3, 596, 598
FES, 574–577, 581, 583
FIB7, 39, 41–42
γ-fibrogen, 207
G6PD, 738, 742
GBSSI, 374–385
GHR, 605, 613–615
H3, 73, 557
HBS1, 598
HMG-CoA, 596, 598
homeobox, 311
HSP70, 596, 598, 600
HSP90, 598
IRBP, 604–614
ITS, 73, 76–77, 298, 336, 370–371, 382
κ-casein, 206–207, 209–212, 215
lactoferrin, 206, 209–212, 215, 218, 220
mitochondrial, 5–6, 10, 16, 32, 41, 74, 439, 451, 460, 462, 510, 572, 587, 649, 651, 656, 674, 677, 680–681, 725
mtLSU-rRNA, 735–738, 740–741
mtSSU-rRNA, 735–738, 740
*NADH*, 460
*ND1*, 73–74, 76, 657, 659
*ND2*, 460, 462–468, 471, 657, 659, 758–759, 762, 777
*ND3*, 657, 659
*ND4*, 161, 163–165, 167, 181, 440, 443, 445–447, 451, 453, 455, 657, 659
*ND4l*, 657, 659
*ND5*, 656–657, 659
*ND6*, 406, 657–659
ndhF, 140
nrDNA, 336
nuclear, 16, 28–29, 32, 41, 74, 375, 386, 572
opsin, 28–30
PcP, 735–744
*Pneumocystis*, 742
protein-coding, 23, 39, 74, 94, 297–298, 307, 358, 396, 405, 445, 460, 471, 506, 604, 609, 612, 656, 682, 705–706
*psaA*, 785, 787, 791
*psaB*, 785, 787–788, 791
RAG-1, 405, 408
RAG-2, 604, 606–610, 613–614
*rbcL*, 140, 336, 379, 696–704
rDNA, 406, 408, 808
RHO1, 574–577, 587, 583
RNA polymerase II, 557
rRNA, 283–284, 297, 307
single-copy, 375
tree, 63, 179, 193, 258, 374, 446, 583–584, 588, 725, 727, 776
tRNA, 2, 52, 94–95, 139, 296–298, 300–301, 303–308, 383, 396, 406, 460, 465–466, 725, 727, 758, 762, 798
tRNAAsp, 674
tRNAL, 73, 76
*trn*L-*trn*F, 380–382, 384
U2 snRNA, 557
v-*myc*, 388
vWF, 605, 614–615, 738
Genes, categories, and species: the evolutionary and cognitive causes of the species problem, Hanno Sandvik (rev.), 565–567
Genetalic
characters, 8, 91
morphology, 71, 82
Genetic analysis, 197
Genomes, 59, 110, 126, 286, 296, 301, 550, 591, 754
chloroplast, 336
mitochondrial, 56, 58–59, 238, 296–297, 301, 303, 307, 309, 462–463, 478, 577, 706
nuclear, 374–375, 571, 604
sequencing, 119, 126
Genomic DNA, 161, 315–316, 443, 550, 566, 574, 606, 758, 823
Genotypes, 735
*Pneumocystis*, 735
Geometric morphometrics, 66–69, 186–205, 488–500
Geometry of the marmot (Rodentia: Sciuridae) mandible: phylogeny and patterns of morphological evolution, Andrea Cardini, 186–205
Geomyidae, 606, 608–609
Geomyoidea, 604–605, 608–611, 613–615
*Geonoma*, 346
*Geotrygon montana*, 38
*Geum*, 374–385
*andicola*, 374, 381–382
*heterocarpum*, 374, 376, 379–384
*montanum*, 374, 376, 379, 382–384
*reptans*, 374, 376, 383–384
*rivale*, 374, 376, 379, 382–384
*schofieldii*, 381
*urbanum*, 374, 376, 379, 382–384
*vernum*, 374, 376, 379, 382–384
Gharials, 386, 398–399
false, 388
true, 388, 396
GHR, 605, 613–615
Giannini, Norberto P., Canonical phylogenetic ordination, 684–695
Gibbon, 52–53
Gillespie, Rosemary G., *see* Hormiga, Gustavo, 70–88
*Ginko*, 784
*Giraffa*, 207, 209, 212, 218, 221–222, 224
*camelopardalis*, 210
Giraffes, 206, 210, 269
Giraffidae, 206–207, 209–210, 212–213, 218–222
Giribet, Gonzalo, Stability in phylogenetic formulations and its relationship to nodal support, 554–564
Gittleman, John L., *see* Bininda-Emonds, Olaf R. P., 724–729
Gliridae, 604–606, 608–609, 611, 614–615
Global clock, 711–712
Gloma fuscipennis, 756
*Glossalictus*, 24
Gnetaleans, 753
Gnetales, 781, 784, 787–788
Goodness-of-fit test, 464, 471, 674
Goose magpie, 388
Gorilla, 52–53, 706–708
Gower, David J., *see* Harris, Simon R., 239–252, 851–852
Gower, David J., *see* Wilkinson, Mark, 127–130
*Gradicrepidula*, 619, 623, 632
*Grantia*, 314–315, 320, 322, 332
*compressa*, 316, 329
Grantiidae, 314, 322–323
*Grantiopsis*, 314–315, 320, 322, 324, 333
sp., 316
Granule-bound starch synthase (GBSSI) (*see also waxy*), 374–385
*Graphiurus murinus*, 606
*Grayia*, 447, 451, 457
*ornata*, 440
Greater grison, 573
Grenyer, Richard, *see* Bininda-Emonds, Olaf R. P., 724–729
Grey-headed chachalaca, 388

Ground spiders, 430
Ground squirrels, 186, 197

*Grus canadensis*, 286
*Gryposuchus colombianus*, 412
Guill, J. Michael, David C. Heins, and Craig S. Hood, Effect of phylogeny on interspecific body shape variation in darters (Pisces: Percidae), 488–500
Guillot, Jacques, *see* Hugot, JeanPierre, 735–744
Guindon, Stéphane and Olivier Gascuel, Simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood, 696–704
Guinea pig, 656–657, 659
*Gulo*, 577, 581, 590
  *gulo*, 573–574
Gymnophthalmidae, 689, 691
Gymnospermae, 518, 524, 781, 787–788

H3, 73, 557
*Haasiella*, 557
Habib, Michael, *see* BinindaEmonds, Olaf R. P., 724–729
*Hadropterus*, 491, 493
Hadziidae, 430
Haemaphysalinae, 299
*Haemaphysalis*, 299, 304–305, 309–310
  *humerosa*, 299
  *inermis*, 299
  *leporispalustris*, 299
  *longicornis*, 299
  ssp., 310
*Haideotriton*, 505, 509
  *wallacei*, 508
Handley, Scott A., *see* Cummings, Michael P., 477–487
*Halictus*, 23–24, 32
  *rubicundus*, 33
Halictidae, 23
Halictid bees, 23–24, 33
Halictini, 24
*Haliotis*, 631
*Halterapis*, 1–2, 4–5, 8–10, 12
  *nigrinervis*, 2, 12
*Hapalemur*, 706
  *griseus*, 736
Haplorhini, 738
Haplotype, 5, 71, 73, 159–160, 165–166, 171–172, 180–181, 465, 467, 776
  tree, 159, 165–166
Hard ticks, 296–298, 300–301, 304, 309
Harris, Simon R., David J. Gower, and Mark Wilkinson, Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), 239–252
Harris, Simon R., David J. Gower, and Mark Wilkinson, Phylogenetic methods and Aetosaur interrelationships: a rejoinder, 851–852
Harshman, John, Christopher J. Huddleston, Jonathan P. Bollback, Thomas J. Parsons, and Michael J. Braun, True and false gharials: a nuclear gene phylogeny of Crocodylia, 386–402
Harvest mouse, 677
Hasegawa-Kishino-Yano (HKY) model, 4, 7, 49, 52–53, 121, 135, 140, 143, 215, 377, 396, 423, 444, 463–464, 466–468, 471, 506, 508, 577, 614, 681–682, 703, 737, 750, 785, 787
Hassanin, Alexandre and Emmanuel J. P. Douzery, Molecular and morphological phylogenies of Ruminantia and the alternative position of the Moschidae, 206–228
Hausdorf, Bernhard and Christian Hennig, Biotic element analysis in biogeography, 717–723
Hawaii, 70–88, 89–109, 820–841
Hawkmoths, 349
Hayashi, Cheryl, *see* Gatesy, John, 403–422
Heckert, Andrew B. and Spencer G. Lucas, Clarifying Aetosaur phylogeny requires more fossils, not more trees—reply to Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), 253–254
Heins, David C., *see* Guill, J. Michael, 4880500
*Helicops*, 447, 451
  *angulatus*, 441
  *pictiventris*, 441
Helminthes, 430, 520
Hemichordata, 296
Hemidactyliine salamanders, 509
Hemihalictus, 23–24, 26, 32–33
*Hemirhagerrhis*, 451
  *nototaenia*, 440
  *viperinus*, 440, 459
Hempel–Oppenheim model, 260, 263
Hendy, M. D., *see* Holland, B. R., 229–238
Hendy, Michael D., *see* Gascuel, Olivier, 110–118
Henicopidae, 559
Henicopinae, 559
Hennig, Christian, *see* Hausdorf, Bernhard, 717–723
Hennig's auxiliary principle, 239, 245–246
Hepaticophyta, 430
*Heptodon*, 857
Herbivores, 339, 349, 746
Herpestidae, 846
*Heteranthera*, 154, 156
  *dubia*, 154, 156
  *limosa*, 154, 156
  *oblongifolia*, 154, 156
  *rotundifolia*, 154, 156
  *seubertiana*, 154, 156
  *zosterifolia*, 154, 156
Heterocoela, 311, 313, 318, 320, 322, 327–328
Heterodactyla, 746, 750–751, 754
*Heterodon*, 447, 451, 455
  *nasicus*, 441
  *simus*, 441
  *platyrhinos*, 287
Heterodonta, 561
Heteromyidae, 606, 608, 614
*Heteropia*, 314–315, 332–333
Heteroptera, 103
Heteroptiidae, 314, 322, 324, 329
*Heterotropus senex*, 756
Heuristic
  parsimony search, 75
  search, 4, 39, 76, 95–96, 166, 318, 327, 336, 368, 372, 377, 504, 575, 607, 609, 611, 621, 666, 737, 823–824, 842, 864
  tree, 607
Heuristics, 368–373, 696, 782, 841
Hexactinellida, 311, 314, 316–318, 320, 328, 330
Hexapoda, 328, 557
Hierarchical phylogenetic models for analyzing multipartitie sequence data, Marc A. Suchard, Christina M. R. Kitchen, Janet S. Sinsheimer, and Robert E. Weiss, 649–664
Hillis, David M., David D. Pollock, Jimmy A. McGuire, and Derrick J. Zwickl, Is sparce taxon sampling a problem for phylogenetic inference?, 124–126
Hillis, David M., *see* Wiens, John J., 501–514
Hippoboscid flies, 37
Hipponicidae, 639
Hipponicids, 620, 622, 632
  artiodactylans, 854

*Hipponix*
sp. Australia, 639
sp. California, 639
Hippopotamidae, 209–210, 221, 855
*Hippopotamus*, 209, 211–213, 215, 222, 706, 712
*amphibius*, 210
Hippos, 207, 210, 707–708, 854–855, 858
*Hippospongia communis*, 316
Hippotragini, 209–210, 215
*Hippotragus*, 209, 212, 215, 218
*niger*, 210
*Hispanomeryx*, 220
Historical
biogeography, 271–280
effects, 684–695
HIV, 660, 662, 735
evolution, 649–664
Hood, Craig S., *see* Guill, J. Michael, 488–500
Holarctic rodents, 186
Holland, B. R., D. Penny, and M. D. Hendy, Outgroup misplacement and phylogenetic inaccuracy under a molecular clock—a simulation study, 229–238
Holothyrida, 300, 309
*Homalictus*, 24
Homalopsinae, 439, 452–453, 455, 457
Homeobox gene, 311
Hominoid, 711–713
Homocoela, 311, 313, 318, 320, 322, 327–328
Homogeneous model, 792
Homoplastic characters, 589, 767
Homoplasy, 16, 30, 55, 57, 61, 152, 154, 156, 211, 230, 239, 243, 245–246, 258, 297, 304, 306–307, 311, 339, 342, 348, 354, 359, 361–362, 398, 403, 415, 420, 442, 444, 460–476, 501–514, 542, 571–572, 575, 577, 581–585, 589, 604, 842, 860–861
index, 762, 764
*Homo*, 706
*sapiens*, 286, 606, 736
*Homoroselaps*, 447, 451, 455–456
*lacteus*, 441
Homoscleromorpha, 328
Honey bees, 26, 296
Honeycreepers, 89
Hood, Craig S., *see* Guill, J. Michael, 488–500
Hoplitomerycidae, 220–221
*Hoplitomeryx*, 220–221
*Hoplobatrachus*
*occipitalis*, 799, 815
*rugulosus*, 799, 815
Horizontal gene transfer (HGT), 649–664
Hormiga, Gustavo, Miquel Arnedo, and Rosemary G. Gillespie, Speciation on a conveyor belt: sequential colonization of the Hawaiian islands by *Orsonwelles* spider (Araneae, Linyphiidae), 70–88
*Hormiphora* sp., 316–317
Horned soldiers, 137, 139–140, 143
Horse flies, 745–746
Horses, 707–708, 857
Horseshoe crab, 300
Host
phylogeny, 740, 742
tree, 737, 740
Host–parasite coevolution, 735–744
House flies, 745
HSP70, 596, 600
HSP90, 598
Huddleston, Christopher J., *see* Harshman, John, 386–402
Huelsenbeck, John P. and Katherine M. Lander, Frequent inconsistency of parsimony under a simple model of cladogenesis, 641–648
Huelsenbeck, John P., Rasmus Nielsen, and Jonathan P. Bollback, Stochastic mapping of morphological characters, 131–158
Hugall, A. F., *see* Lee, M. S. Y., 15–22
Hugot, Jean-Pierre, Christine Demanche, Véronique Barriel, Eduardo Dei-Cas, and Jacques Guillot, Phylogenetic systematics and evolution of primate-derived *Pneumocystis* based on mitochondrial or nuclear DNA sequence comparison, 735–744
Human, 52–53, 526, 565, 571, 591, 656707–708, 711–713, 715, 738
Humphries, Christopher J., *see* Williams, David M., 255–258
Hyaenidae, 846
*Hyalomma*, 299, 304–305, 310
*aegyptium*, 299
*dromedarii*, 299
*marginatum*, 299
*truncatum*, 299
Hybridization, 63, 89, 92, 103, 379, 383, 820, 832
*Hydrodynastes*, 447, 451
*bicinctus*, 441
Hydrophiinae, 441, 447
*Hydrophis*, 447, 451, 456
*semperi*, 441
*Hydropotes*, 206, 209, 212, 215, 218–220, 222, 224–225
*inermis*, 210
Hydropotinae, 209
Hydropsini, 456
*Hydrothrix gardneri*, 154, 156
*Hyemoschus*, 206, 219
*Hylaeochampsa*, 412
Hymenoptera, 23, 296, 307
Hypertragulidae, 207, 222
*Hypertragulus*, 221
*Hyposmocoma*, 89
Hypothesis test, 48–54, 781–793
*Hyracotherium*, 857
Hystricognathi, 604–605, 608–611, 613, 615

*Ictonyx*, 590
*striatus*, 581, 589–590
Identifying conflicting signal in a multigene analysis reveals a highly resolved tree: the phylogeny of Rodentia (Mammalia), Ronald W. DeBry, 604–617
Idiosyncratic markers, 296–310
Impala, 210
Ichthyosaurs, 501
Incomplete taxa, 528–538
Incongruence, 11, 15, 20, 28, 37, 39–41, 43, 45, 404, 451, 559, 571–572, 581, 587, 608, 613
length difference, 75–77, 406, 408, 463
length difference (ILD) test (*see also* partition homogeneity test), 3, 11, 15, 26, 28, 75–76, 418, 467, 604, 607–609, 613, 622, 773, 853–855, 860, 862
Inconsistency, 229–238, 641–648
Incorrect model, 674–683
Indel-change ratio, 556–557
Indels, 283–295, 300, 377–378, 392, 394, 397, 574–575, 651, 823, 827
characters, 389, 823, 826, 829
coding, 444, 575
parsimony informative, 823
Independent and identically distributed (IID), 842, 847
Ingroup, 27, 56, 89, 91–92, 96–97, 99, 197, 229–230, 386–387, 389, 392, 394, 396–397, 443, 576, 620–622, 748, 750, 753, 757, 764, 775, 824, 826

Informative characters, 16, 575, 583
*Inquilina*, 1
*dawsoni*, 2
*schwarzi*, 2
Insecta, 37, 296, 430
Insects, 23, 32, 34, 89, 106, 430
Insertion (*see* indel)
Internal transcriber spacer (ITS), 73, 76–77, 298, 336, 370–371, 382, 820–841
Intraorganismal homology, character construction, and the phylogeny of Aetosaurian Archosaurs (Reptilia, Diapsida), Simon R. Harris, David J. Gower, and Mark Wilkinson, 239–252
Intraspecific phylogeny of the New Zealand short–tailed bat Mystacina tuberculata inferred from multiple mitochondrial gene sequences, Brian D. Lloyd, 460–476
Intron–exon, 20, 26
Introns, 376, 383, 391, 394, 571–593
Invertebrates, 429, 618–619
*Ips* beetles, 677
IRBP, 604–614, 738
Irregularia, 328
*Ischnura*, 91, 94, 102
*aurora*, 94
*heterosticta*, 94
Island evolution, 820–841
Iskandar, Djoko, *see* Evans, Ben J., 794–823
Isopoda, 430
Is sparce taxon sampling a problem for phylogenetic inference?, David M. Hillis, David D. Pollock, Jimmy A. McGuire, and Derrick J. Zwickl, 124–126
*Ixodes*
*affinis*, 300
*auritulus*, 300
*hexagonus*, 299–300
*holocyclus*, 300
*pilosus*, 300
*simplex*, 300
ss., 298, 300, 304, 307, 309
*tasmani*, 300, 304, 309
*uriae*, 300
Ixodidae, 296–310

Jacarea, 409, 412
Jaccard distance, 718
Jackknife, 55–65, 128–129, 409, 554, 556, 560–561, 575, 581
analysis, 407
parsimony, 129, 557, 559
scores, 412
support, 419, 540, 559, 561, 582–583
values, 589, 665
*Janacus*, 619, 623, 632, 636, 639
*Janneria pustulata*, 640
Japygidae, 557
Jean-Marie, Alain, *see* Gascuel, Olivier, 110–118
*Jenkina*, 314–315
Jenkinidae, 314
Jennings, W. Bryan, Eric R. Pianka, and Stephen Donnellan, Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas, 757–780
Ji, Shuqing, *see* Danforth, Bryan N., 23–36
Johnson, Kevin P., Richard J. Adams, Roderic D. M. Page, and Dale H. Clayton, When do parasites fail to speciate in response to host speciation?, 37–47
Jones, Kate E., *see* Bininda-Emonds, Olaf R. P., 724–729
Jordan, Steve, Chris Simon, and Dan Polhemus, Molecular systematics and adaptive radiation of Hawaii's endemic damselfly genus *Megalagrion* (Odonata: Coenagrionidae), 89–109
Joyce, Paul, *see* Minin, Vladimir, 674–683
Jukes–Cantor
distances, 425
model, 133, 371, 424, 485, 530, 644, 666–669, 703, 708, 711–715, 784–785, 787–788, 820, 825, 831–832, 834
Juliformia, 557
Jungles
analysis, 39, 43, 45
optimizations, 42
Jurassic, 746, 753
late-, 549, 746
lower, 746, 749
mid-, 745
upper, 753

$\kappa$-casein, 206–207, 209–212, 215
*Kebira*, 333
Kelly, Christopher M., R. Nigel P. Barker, and Martin H. Villet, Phylogenetics of advanced snakes (Caenophidia) based on four mitochondrial genes, 439–459
Kendall's
shape space, 66
tangent space, 66–67
Kimura three-parameter model, 577, 581
Kimura two-parameter model, 133, 229, 231, 236, 485, 577, 700, 703, 762, 801, 823, 826, 850
Kishino-Hasegawa test, 4, 10, 15–22, 212, 216, 390, 392, 394–396, 398, 445, 504, 781
Kishino, Hirohisa, *see* Wiegmann, Brian M., 745–756
Kitchen, Christina M. R., *see* Suchard, Marc A., 649–664
Kiwi, 474, 550
Kjer, Karl M., *see* Xia, Xuhua, 283–295
Koepfli Klaus-Peter and Robert K. Wayne, Type I STS markers are more informative than cytochrome *b* in phylogenetic reconstruction of the Mustelidae (Mammalia: Carnivora), 571–593
Kolmogorov–Smirnov test, 190
*Krebsia* sp., 639
Kulczynski distance, 718
Kullback–Leibler distance, 782, 785
Kumar, Sudhir, *see* Rosenberg, Michael S., 119–123

*Labulla*, 71, 77
*thoracica*, 71–73, 88
Lacanicephalidea, 430
Lactoferrin, 206, 209–212, 215, 218, 220
*Lagomeryx*, 224
Lagomorpha, 606, 839–851
*Laminacauda gigas*, 71
*Lamontia*, 314–315, 317, 323–324
*zona*, 333
Lamprophiinae, 439, 441, 447, 455–456
*Lamprophis*
*fuliginosus*, 441
*inornatus*, 441, 445, 459
Lander, Katherine M., *see* Huelsenbeck, John P., 641–648
*Lanice conchylega*, 316
Lapointe, François-Joseph, Mark Wilkinson, and David Bryant, Matrix representations with parsimony or with distances: two sides of the same coin?, 865–868
Lapointe, François-Joseph, *see* Wilkinson, Mark, 127–130
*Larus glaucoides*, 286
Larvae, 8, 618, 631
Larval characters, 8–9
*Lasioglossum*, 23–28, 30, 32–33

(Dialictus) cressonii, 27
(Dialictus) figueresi, 27, 33
(Dialictus) gundlachii, 27
(Dialictus) hyalinum, 27
(Dialictus) imitatum, 27
(Dialictus) parvum, 27
(Dialictus) pilosum, 27
(Dialictus) rohweri, 27
(Dialictus) tegulare, 27
(Dialictus) umbripenne, 27
(Dialictus) vierecki, 27
(Dialictus) zephyrum, 24, 27, 33
(Evylaeus) albipes, 26–27, 33
(Evylaeus) apristum, 27
(Evylaeus) boreale, 27
(Evylaeus) brevicorne, 27
(Evylaeus) calceatum, 24, 27, 33
(Evylaeus) cinctipes, 27
(Evylaeus) comagenense, 27
(Evylaeus) duplex, 27
(Evylaeus) fulvicorne, 27
(Evylaeus) gattaca, 27
(Evylaeus) inconditum, 27
(Evylaeus) interruptum, 27
(Evylaeus) laticeps, 24, 27
(Evylaeus) limbellum, 27
(Evylaeus) lineare, 27
(Evylaeus) lucidulum, 27
(Evylaeus) malachurum, 27
(Evylaeus) marginatum, 24, 27, 33
(Evylaeus) mediterraneum, 27
(Evylaeus) morio, 27
(Evylaeus) nigripes, 27
(Evylaeus) pauxillum, 27
(Evylaeus) pectorale, 27
(Evylaeus) politum, 24, 27, 33
(Evylaeus) puncticolle, 27
(Evylaeus) quebecense, 27
(Evylaeus) subtropicum, 27
(Evylaeus) truncatum, 27
(Evylaeus) villosulum, 26–27
(Hemihalictus) lustrans, 24, 27, 33
(Lasioglossum) athabascense, 27
(Lasioglossum) coriaceum, 26
(Lasioglossum) fuscipenne, 27
(Lasioglossum)leucozonium, 24
(Lasioglossum) pavonotum, 27
(Lasioglossum) sisymbrii, 27
(Lasioglossum) titusi, 27
(Paralictus) asteris, 27
(Sphecodogastra) noctivaga, 27
(Sphecodogastra) oenotherae, 27
(Sudila) alphenum, 27
*Laticauda*, 451
  *colubrina*, 441
*Latimeria chalumnae*, 286
*Laupala*, 833
Least and most powerful phylogenetic tests to elucidate the origin of the seed plants in the presence of conflicting signals under misspecified models, Stéphane Aris–Brosou, 781–793
Least-squares
  algorithm, 866
  criterion, 865
Lee, M. S. Y. and A. F. Hugall, Partitioned likelihood support and the evaluation of data set conflict, 15–22
Leeches, 430
*Leioheterodon*, 451, 456
  *madagascariensis*, 441
*Lelapia*, 314–315, 324
Lelapiidae, 314, 329
Lemuridae, 711–713
Lemuriform, 711–713
Lemurs, 52–53, 524, 706,
  mouse, 705–706, 708, 711–715
Le Parco, Yannick, *see* Manuel, Michael, 311–333
*Lepilemur*, 706
*Lepthyphantes minutus*, 73, 88
*Leptictis*, 858
Leptomerycidae, 222
*Leptomeryx*, 222
*Leptonetis perplexus*, 639
*Leptonycteris curasoae yerbabuenae*, 473
*Leptotila*, 41
  *jamaicensis*, 38, 41–42
  *plumbeiceps*, 38
  *rufaxilla*, 38
  *verreauxi*, 41
  *verreauxi angelica*, 38
  *verreauxi fulviventris*, 38
*Leptynoma hessei*, 756
*Lestis*, 2
Leucaltidae, 314
*Leucaltis*, 314–315, 320, 324, 332
  *clathria*, 316
*Leucandra*, 314–315, 320, 322–324, 332–333
  *aspera*, 316
*Leucascandra*, 314–315, 332
Leucascidae, 314
*Leucetta*, 314–315, 320, 322, 324, 330, 332–333
  *chagosensis*, 316
Leucettidae, 314
*Leucilla*, 314–315, 332
*Leucocrossuromys leucurus*, 189
*Leuconia*, 314–315, 320, 322–324, 328, 330, 332–333
  *johnstoni*, 333
  *nivea*, 316, 329, 333
  *ochotensis*, 333
*Leucopsila*, 314–315, 317, 323
*Leucosolenia*, 314–315, 320, 322–323, 329, 332–333
  *laxa*, 332
  sp., 316
  *variabilis*, 329
Leucosoleniida, 314, 320, 323
Leucosoleniidae, 314
Levin, Rachel A., Lucinda A. McDade, and Robert A. Raguso, Systematic utility of floral and vegetative fragrance in two genera of Nyctaginaceae, 334–351
*Lialis*, 757–758, 762, 764, 767, 770–771, 775
  *burtonis*, 759, 762
  *jicari*, 759
Liancalus sp., 756
Lice, 37–47, 430
Lichens, 430, 518
*Ligeromeryx*, 224
Likelihood (see maximum likelihood), 352–367
  analysis, 18, 263, 374, 708, 715, 803–804, 807
  framework, 369
  methods, 286
  model, 17, 212, 356–358, 362, 481, 506, 555, 674, 681, 705
  ratio tests (LRT), 41–42, 48, 54, 63, 95, 352, 363–364, 377, 379, 390, 394, 445, 575, 580, 598, 600, 604–617, 621, 655, 674–683, 714, 825, 849, 801

trees, 39, 506, 508, 762, 764, 802
weight test, 18–19, 22
*Limnonectes*, 794–795, 797–799, 802, 805–809, 811–812, 815
*acanthi*, 797, 802, 807, 809–811, 815
*arathooni*, 795, 797, 812, 815
*asperatus*, 797, 799, 815
*blythii*, 797, 808
cf. *acanthi*, 802, 810, 815
cf. *blythii* 1, 804, 815
cf. *blythii* 2, 804, 815
cf. *blythii* 3, 802, 804, 815
cf. *heinrichi*, 812, 815
cf. *kuhlii* 1, 795, 797, 808, 815
cf. *kuhlii* 2, 795, 797, 808, 815
cf. *kuhlii* 3, 795, 797, 808, 815
cf. *magnus*, 797, 802, 807, 811, 815–16
cf. *microtympanum* 1, 795, 797, 802, 812, 816
cf. *microtympanum* 2, 795, 797, 802, 812, 816
cf. *modestus*, 795, 797, 802, 812, 816
cf. *modestus* complex 2, 797, 802, 812, 816
*dammermani*, 809
*diuatus*, 811
*finchi*, 802, 804, 816
*grunniens*, 795, 802, 804, 809, 816
*gyldenstolpei*, 797, 816
*heinrichi*, 795, 797, 816
*ibanorum*, 802, 804, 816
*ingeri*, 797, 804, 816
*kardasani*, 809, 816
*laticeps*, 797, 816
*leporinus*, 797, 816
*leytensis*, 797, 802, 807, 809–812, 816–817
*limborgii*, 797
*macrocephalus*, 797, 802, 807, 811, 817
*macrodon*, 797, 804, 816, 8167
*magnus*, 797, 802, 807, 811, 817
*malesianus*, 797, 800, 804, 808, 816, 817
*micrixalus*, 811
*microdiscus*, 795, 809, 817
*palavanensis*, 802, 804, 811, 817
*paramacrodon*, 802, 804, 808, 817
*parvus*, 802, 804–807, 811, 817
*pileata*, 797
*shompenorum*, 797
sp. 1, 797, 817
sp. 2, 797, 817
sp. D, 797, 812, 817
sp. G, 817–818
sp. G2, 797, 812
sp. I, 802, 804, 807, 812, 818
sp. nov., 797
sp. T, 797, 812, 818
sp. V complex 1, 797, 812, 818
sp. V complex 2, 797, 802, 819
*visayanus*, 797, 802, 807, 811, 819
*woodworthi*, 797, 802, 807, 811, 819
*Limulus polyphemus*, 300
Linear model, 684–695
Linnear parsimony, 342
*Linyphia*, 77
*triangularis*, 73, 88
Linyphiidae, 70–88
Linyphiini, 72, 77
*Liocheles waigiensis*, 300
Liolaemidae, 159
*Liolaemus*, 159–185, 688
*austromendocinus*, 161, 163, 167, 178–179
*bibronii*, 164, 167
*buergeri*, 161, 178, 182
*capillitas*, 161, 163, 167, 179
*ceii*, 161, 178, 182
*ceii–kriegi* clade, 172
*christiani*, 161
*curis*, 161
*elongatus*, 159–161, 163, 166–167, 171–172, 178–179, 182
*elongatus–kriegi* complex, 159–161, 167, 182
*gracilis*, 164, 167
*heliodermis*, 161
*kingii*, 164, 167
*kriegi*, 159–161, 164, 166–167, 171–172, 178, 182
*leopardinus*, 161
*lineomaculatus*, 164, 167
*neuquensis*, 164, 167
*petrophilus*, 161, 163, 166–167, 172, 178–179, 182
*pictus*, 164, 167
*pseudocnomalus*, 165, 167
sp. 1, 163, 182
sp. 2, 164, 179
sp. 3, 163–164, 167
sp. 4, 164, 179, 182
sp. 5, 163, 171, 178, 182
sp. 6, 163, 171, 178, 182
sp. 7, 163, 171, 178, 182
sp. 8, 164, 179, 182
sp. A, 164, 179, 182
sp. B, 164, 178, 182
sp. C, 164, 178
*thermarum*, 161, 178
*vallecurensis*, 164, 167
*Liophis*, 447, 451
*reginae*, 441
Lithonida, 314, 317, 323, 329–330
*Littorina littorea*, 640
Liverworts, 430
Lizards, 105, 159–185, 286, 306, 423, 689, 691, 693, 757–780
agamid, 758
New World, 160
pygopodid, 757–780
Llamas, 221
Lloyd, Brian D., Intraspecific phylogeny of the New Zealand short–tailed bat Mystacina tuberculata inferred from multiple mitochondrial gene sequences, 460–476
Lobelioids, 833
Local clock analysis, 89
Local molecular clock, 89–109, 705–716
LogDet
analysis, 3, 6–7, 11
method, 784
tree, 4, 7
LogDet/Paralinear
analysis, 614, 621
distance (*see also* paralinear distance), 284, 287, 289, 291, 293, 397, 580, 610, 612, 676
transformation, 579
Log-likelihood, 596, 598–599, 614, 655, 782
ratio test, 3–4
scores, 445, 464, 607, 761
values, 445, 761
Long branch
attraction, 63, 79, 82, 96, 327, 354–370, 386, 419, 423–428, 443, 478, 483, 510, 531, 537, 558, 590, 594
effects, 2

subdivision, 423–428
Longhorn beetle, 89
Longirostres, 390–391, 398, 400
*Longosuchus*, 241–243, 247–252
Long wavelength (LW)
opsin, 26–28
rhodopsin, 23
*Lontra*, 577, 590
*canadensis*, 573
*felina*, 573
*longicaudis*, 573
*Lorancameryx*, 224
*Lordotus* sp., 756
*Loris*, 706
Lorisiform, 711–713
Loxocemidae, 440
*Loxocemus bicolour*, 440
*Loxodonta*, 591
Lucas, Spencer G., *see* Heckert, Andrew B., 253–255
*Lucasuchus*, 241–243, 250–251
*Lutra*, 590
*lutra*, 573
*maculicollis*, 583, 590
Lutrinae, 571–573, 581, 590, 846
*Lycodon*, 447, 451, 456–457
*capucinus*, 440
*laoensis*, 440
*Lycodonomorphus*, 456
Lycodontinae, 456
Lycodontini, 457

Macaca, 706, 741
*fascicularis*, 736, 741
*mulatta*, 736
*nemestrina*, 736, 741
Macaque, 52–53
*Macroderma gigas*, 472
*Macrogalea*, 1, 4, 7–8, 10–13
*candida*, 2, 10, 12
*scaevolae*, 12
*zanzibarica*, 2, 10, 12
*Macropisthodon*, 447, 451, 457
*rudis*, 440, 447, 456
Macrostomata, 440
*Madagascarophis*, 451
*colubrine*, 441
Magnoliaceae, 346
Magnolias, 346
Magpie goose, 388
Mahalanobis distance, 192, 491
Malacostraca, 557
*Malluvium* sp., 639
Maloideae, 375
*Malpolon*, 451
*monspessulanus*, 441, 459
Mammalia, 566, 571–593, 604–605, 861
Mammalian phylogeny, 857
Mammals, 16, 23, 56, 66, 106, 206, 219, 224, 261–262, 283–287, 293, 415, 518, 524, 550, 571, 605, 794, 853, 861
eutherian, 249, 585, 860
placental, 591
Mandible, 186–205
Mandibular
characters, 192
dendograms, 195
Mandibulata, 557
*Manuelia*, 8
*postica*, 2
Manueliini, 1, 9
Mantel
matrix correlations, 493–494, 497
statistics, 488, 491, 493
test, 491
Manuel, Michael, Carole Borchiellini, Eliane Alivon, Yannick Le Parco, Jean Vacelet, and Nicole Boury-Esnault, Phylogeny and evolution of Calcareous sponges: monophyly of Calcinea and Calcaronea, high level of morphological homoplasy, and the primitive nature of axial symmetry, 311–333
*Maoricicada*, 674
*campbelli*, 474
*Maoricrypta*, 619, 623, 631–632, 636
Markov chain (*see also* MCMC), 50, 133–134, 140, 165, 212, 377, 445, 576, 666, 749–750, 760, 785
continuous time, 179, 644
model, 782, 785, 791, 841
processes, 594
Markov chain Monte Carlo (MCMC), 48–54, 95, 131–158, 159, 212, 377, 445, 478, 484, 576, 604, 607, 649–664, 667, 674, 709, 711, 749, 751, 760
algorithms, 135–136, 665–666, 697, 711, 785
analysis, 140, 213, 486, 750, 801
approach, 702, 750
Metropolis-coupled, 604, 607, 621, 623
procedure, 748, 750
processes, 825
sampling, 477–487, 612
simulations, 165
*Marmota*, 186–205
*baibacina*, 189, 192–193, 196, 199, 201
*baibacina centralis*, 201
*bobak*, 189, 192, 196–197, 199, 202
*broweri*, 186, 189–190, 192–193, 196–198, 202
*caligata*, 186, 190, 192, 194–196, 198, 202
*caligata caligata*, 202
*caligata cascadensis*, 202
*caligata nivaria*, 202
*caligata okanagana*, 202
*caligata oxytona*, 202
*caligata sheldoni*, 202
*caligata vigilis*, 202
*camtschatica*, 186, 189, 192, 196, 198
*camtschatica bungei*, 202
*caudata*, 188–190, 192, 195–196, 199
*caudata aurea*, 202
*flaviventris*, 192, 194–197, 205
*flaviventris avara*, 203
*flaviventris dacota*, 203
*flaviventris engelhardti*, 203
*flaviventris flaviventris*, 203
*flaviventris luteola*, 203
*flaviventris nosphora*, 203
*flaviventris obscura*, 203
*flaviventris parvula*, 203
*flaviventris sierrae*, 203
*himalayana*, 188–189, 195–197, 203
*himalayana himalayana*, 203
*himalayana robusta*, 203
*marmota*, 186–187, 189, 192, 196, 203
*marmota marmota*, 203–204
*menzbieri*, 188–189, 192, 195–197, 199, 204
*monax*, 186, 189, 192–193, 196–197, 199, 204
*monax canadensis*, 204
*monax ignava*, 204
*monax monax*, 204

*monax ochracea*, 204
*monax petrensis*, 204
*monax preblorum*, 204
*monax rufescens*, 204
*olympus*, 186, 190, 192–198, 204
*sibirica*, 189–190, 192–193, 196, 199, 204
*vancouverensis*, 186, 188, 190, 192, 194, 196, 198, 204
Marmotini, 197
Marmots (*see also Marmota*), 186, 188–189, 192–195, 197, 205
Alaskan, 186, 193, 195, 198
Alpine, 186–187, 198–199
black-capped, 186, 193, 198–199
Eurasiatic, 186, 193, 198–199
hoary, 194
long-tailed, 195, 199
Menzbier, 189, 199
Olympic, 194–195, 197–198
Palaearctic, 187, 198–199
Vancouver Island, 189, 194, 196–199
yellow-bellied, 196–198, 205
Marsupials, 524
Martens, 571–573
American, 573
*Martes*, 577, 581, 590
*americana*, 573, 581, 590
*pennanti*, 573, 581, 590
Marsupials, 709
Mast, Austin R. and Reto Nyffeler, Using a null model to recognize significant co-occurrence prior to identifying candidate areas of endemism, 271–280
*Maticora*, 447, 451
*bivigata*, 441
Matrix correlation analysis,
nesting, 241
parsimony, 241
with distances (MRD), 865–868
with parsimony (MRP), 255, 439, 455–456, 839–851, 865–868
Matrix representations with parsimony or with distances: two sides of the same coin? François-Joseph Lapointe, Mark Wilkinson, and David Bryant, 865–868
Matrix-weighted
analysis, 622
parsimony, 623
Maximum likelihood, 16–18, 23, 26, 28, 39, 41, 63, 76, 95, 98–100, 121, 128, 133, 159, 229–238, 283, 285, 287, 293, 320, 363, 365, 368–373, 386, 439–459, 463, 468, 477–487, 506, 571, 585, 594–603, 604–617, 622, 641–642, 656, 665–666, 671–683, 696–704, 705–716, 726, 801
analysis, 1, 3, 7, 11, 26–27, 30, 42, 63, 76, 79, 89, 95–97, 99, 166–167, 171, 182, 211–212, 215, 231, 318, 322, 327, 329, 341, 377–378, 386, 389–90, 394–396, 407, 419, 715, 737, 740, 797, 801, 824, 829, 850
bootstrap support, 667–668
estimates, 166, 394, 594, 596, 601, 604, 669, 709, 711–712, 714, 748, 850
model, 11, 45, 96, 159, 220, 358, 397
searches, 95
tests, 614
trees, 12, 18, 20, 22, 30, 42–45, 81, 96, 128, 323, 328, 368, 371, 383, 396, 455–456, 479, 574, 576–577, 580–581, 583, 589–590, 607, 611, 666, 677, 681, 762, 767, 771, 781, 825, 827–829, 849
value, 140
Maximum parsimony (*see also* parsimony), 1, 12, 95, 98, 120, 159, 229–238, 293, 298, 306–307, 318, 327, 339–340, 425, 439–459, 463, 468, 478, 489, 607, 609–610, 613–614, 726, 801, 820, 823, 826, 833
analysis, 3, 5–7, 10–11, 95, 166–167, 182, 211, 213, 215, 275, 317, 322, 327, 447, 467, 571, 577, 580–581, 615, 737, 740, 797, 801, 824, 829, 850
reconstruction, 450
trees (*see also* most-parsimonious trees), 4, 110, 213, 215, 217, 240, 242–243, 248, 252, 275, 317, 320, 323, 329, 336, 339–341, 346, 359, 378, 383–384, 444, 446–447, 463, 574, 576–577, 579–581, 589–590, 607–608, 824, 826–828
McDade, Lucinda A., *see* Levin, Rachel A., 334–351
McGuire, Jimmy A., *see* Evans, Ben J., 794–823
McGuire, Jimmy A., *see* Hillis, David M., 124–126
McLachlan, Robert, *see* Gascuel, Olivier, 110–118
*Megalagrion*, 89–109, 833
*adytum*, 93, 103, 105
*blackburni*, 93, 103
*calliphya*, 93, 103
damselflies, 89–109, 833
*eudytum*, 89, 93, 103, 105
*hawaiiense*, 93, 101–104, 106
*heterogamias*, 93, 101, 103–104, 106
*jugorum*, 103, 106
*kauaiense*, 93, 101, 103–104, 106
*koelense*, 92–93, 101, 104
*leptodemas*, 93, 103
*mauka*, 89, 93, 101, 103, 105
*molokaiense*, 106
*nesiotes*, 89, 93, 101, 103, 105–106
*nigrohamatum*, 93, 103
*nigrolineatum*, 93
*oahuense*, 89, 93, 101, 103, 105
*oceanicum*, 93, 103
*oresitrophum*, 89, 93, 99, 101, 103–104, 106
*orobates*, 89, 93, 99, 101, 103–104, 106
*pacificum*, 89, 92–93, 99–100, 103, 105–106
*paludicola*, 89, 93, 101, 103, 105–106
sp. nov., 93
*vagabundum*, 89, 93, 103, 105
*williamsoni*, 92, 103
*xanthomelas*, 89, 91–93, 103, 105
Megaloptera, 106
Megapodon, 314–315, 328, 333
Megophryidae, 794
*Mehelya*, 447, 451, 455–456
*capensis*, 441
*nyassae*, 441, 459
*Melanesobasis*, 91
*Melanosuchus*, 391, 396, 412
*niger*, 388, 405
*Meles*, 581, 590
*meles*, 573–574, 577, 582
Melinae, 571–573, 581
Mellivorinae, 572–573
Melnick, Don J., *see* Evans, Ben J., 794–823
*Melogale*, 589–590
*moschata*, 581–583, 590
Mephitinae, 572–573, 590, 846
Merostostomata, 300
Merycodontinae, 224
Mesonychia, 853–855, 858, 860
Mesostigmata, 300, 309
Mesozoic, 549–553, 745–746
Meta-optimality criterion, 554–564
Metastriata, 296, 299, 304, 307
Metazoa, 311–333, 520
Meteoriaceae, 430
*Methanococcus jannaschii*, 659
*Metriopelia ceciliae*, 38
Metropolis-coupled MCMC, 604, 607, 621, 623
Metropolis–Hastings

algorithm, 662
criterion, 50
proposal, 654–655
Metropolis-within-Gibbs cycle, 654–655
*Metrosideros polymorpha*, 820
Microbats, 460
Microbes, 429
*Microcebus*
*berthae*, 706
*griseorufus*, 706
*murinus*, 706
*myoxinus*, 706
*ravelobensis*, 706
*rufus*, 706
*sambiranensis*, 706
*tavaratra*, 706
Microchiroptera, 460–476
*Microciona prolifera*, 316
Microevolutionary model, 692
Microhylidae, 794
*Microlinyphia*, 77
*dana*, 73, 88
*Micruroides*, 447, 451
*euryxanthus*, 441
*Micrurus*, 447, 451
*diastema*, 441
*fulvius*, 441
Miller, Jeremy A., Assessing progress in systematics with continuous jackknife function analysis, 55–65
Millipedes, 430
*Minchinella*, 314–315, 317, 323, 328–329, 332
Minchinellidae, 314, 320, 328
Minimum evolution, 121, 287, 478, 489, 579
tree, 506, 580, 610
Minimum-length trees, 415, 750
Minin, Vladimir, Zaid Abdo, Paul Joyce, and Jack Sullivan, Performance-based selection of likelihood models for phylogeny estimation, 674–683
Mink, American, 573
Miocene, 207, 221, 224–225, 451, 549, 629, 761, 776
early-, 206–207, 209, 216, 218, 224–225
mid-, 206, 217–218, 224–225
late-, 218, 224
*Mirabilis*, 334–351
*alipes*, 340
*bigelovii*, 336
*greenei*, 343–344
*jalapa*, 336
*longiflora*, 336
*macfarlanei*, 341, 343–344
*multiflora*, 347
*pudica*, 340
*Mirza*, 706
Misleading zones, 229–238
Missing data, 528–538
Missing data, incomplete taxa, and phylogenetic accuracy, John J. Wiens, 528–538
Mites, 299–300, 430
Mitochondrial
alleles, 41
DNA, 3–4, 7, 12, 37, 48, 52, 89, 94–101, 103, 159–185, 296, 386–387, 395, 407–408, 415, 418, 423, 439–459, 460–461, 464, 471, 474, 503–504, 506, 508–509, 548, 551, 571–572, 583, 586, 589, 656, 674, 677, 680–681, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776, 794, 798, 801, 808, 810–812
DNA tree, 178
genes, 5–6, 10, 16, 32, 41, 74, 439, 451, 460, 462, 510, 572, 587, 649, 651, 656
genomes, 56, 58–59, 238, 296–297, 301, 303, 307, 309, 462–463, 478, 577, 706
RNA, 445
rRNA, 615
tRNA, 615
*Mnemiopsis*, 317
*leidyi*, 316
Models
analysis, 542
Bayesian, 49, 650
birth–death, 641, 644–645, 647, 761
cladogenetic, 641
continuous-time Markov, 131–134, 147, 149, 651
discrete, 594, 602
DNA substitution, 444
equal-rates, 57
evolutionary, 126, 352, 652, 665, 668, 672, 692, 696–697, 705, 727, 781–782, 788, 791–792, 801, 829, 850
Felsenstein, 463, 703, 708–709, 711–715, 748, 850
gamma, 594–603, 750
general-time-reversible (GTR), 3–4, 7–8, 18–19, 26, 28–30, 42, 95, 97, 165–166, 212, 286, 318, 390, 392, 396, 445, 463, 576, 579, 585, 608–611, 613, 666–669, 674, 677, 679, 681–682, 737, 762, 784, 801–802, 825, 831, 834
Hasegawa-Kishino-Yano (HKY) , 4, 7, 49, 52–53, 121, 135, 140, 143, 215, 377, 396, 423, 444, 463–464, 466–468, 471, 506, 508, 577, 614, 681–682, 703, 737, 750, 785, 787
Hempel–Oppenheim, 260, 263
homogeneous, 792
incorrect, 674–683
Jukes–Cantor, 133, 371, 424, 485, 530, 644, 666–669, 703, 708, 711–715, 784–785, 787–788, 820, 825, 831–832, 834
Kimura three-parameter, 577
Kimura two-parameter, 133, 229, 231, 236, 485, 577, 700, 703, 762, 801, 823, 826, 850
likelihood, 17, 212, 356–358, 362, 481, 506, 555, 674, 681, 705
linear, 684–695
Markov, 147, 594–603, 782, 785, 791, 841
maximum likelihood, 11, 45, 96, 159, 220, 358, 397
microevolutionary, 692
misspecification, 665–673
molecular clock, 48
nested, 606
nonreversible nucleotide substitution, 48
nuclear substitution, 215, 287, 377, 604–617, 658, 674–683, 708, 785
null, 271–280, 480, 482, 484, 717–723
parsimony, 361, 363
phylogenetic, 133, 594, 649, 651, 654, 656, 661
Poisson process, 134
rates-across-sites, 595–596
rate homogeneity, 463
regression, 684, 686
site-specific-rates (SSR), 4, 8, 26–28, 30, 463, 466–468, 610, 612, 674
stochastic, 152, 705, 761, 833
substitution, 3, 140, 293, 371, 463, 577
Tamura and Nei, 703
three-parameter TN93, 293
time-homogeneous, 647
tree, 231, 423, 427, 540, 542, 761, 841–842
Wright–Fisher, 179
Model-based Gaussian clustering (MBGC), 717–718, 722
Models of evolution, 126, 352, 364, 377, 615, 665, 668, 672, 692, 696–697, 705, 727, 781–782, 788, 791–792, 801, 829, 850
Molecular

analysis, 103–104, 221, 323, 386, 397, 796
characters, 55, 70, 76–77, 131, 199, 209, 296, 298
cladograms, 187, 742
cladistics, 187
clock (*see also* Bayesian clocklike model, non–parametric rate smoothing), 11, 48, 50–53, 70, 81, 110, 229–231, 233, 386, 390, 395, 550, 641, 644, 647, 700, 705–716, 745, 754, 757–780, 782, 801–802, 825, 841
data, 57, 62, 75, 82, 99, 103–105, 187, 300, 317, 320, 328–329, 339, 369, 386–389, 397, 401, 403, 406, 423, 502–504, 510–511, 539, 543, 726–727, 760, 764, 767, 770, 796, 853, 857
dating analysis, 375
evolution, 20, 38, 41, 591, 594–603, 785, 791
local, 89–109
model, 48
partitions, 17
phylogenetics, 187, 283–295, 311, 715
phylogeny, 37–47, 374–385, 545, 781, 832
studies, 206, 386, 501, 504, 604, 614, 727
substitution, 37
systematics, 430, 501–514, 571–572, 591, 745–756, 820–841
tree, 103, 105–106, 386, 389–392, 394–396, 399, 403, 853, 858, 860
Molecular and morphological phylogenies of Ruminantia and the alternative position of the Moschidae, Alexandre Hassanin and Emmanuel J. P. Douzery, 206–228
Molecular phylogenetics of Allodapine bees, with implications for the evolution of sociality and progressive rearing, Michael P. Schwarz, Nicholas J. Bull, and Steven J. B. Cooper, 1–14
Molecular systematics and adaptive radiation of Hawaii's endemic damselfly genus *Megalagrion* (Odonata: Coenagrionidae), Steve Jordan, Chris Simon, and Dan Polhemus, 89–109
Molecules, 206–228
Mollusks, 430, 520, 619, 633
deep sea, 430
Monkeys, 707–708, 811
owl, 738
Monochoria
*cyanea*, 154, 156
*hastata*, 154, 156
*korsakovii*, 154, 156
*vaginalis*, 154, 156
Monocotyledones, 524
*Monoplectroninia*, 329
*Monosiga brevicolis*, 316
Monte Carlo
error, 750
estimate, 50
permutation, 684–695
randomization test, 684
simulation (*see also* computer simulation and simulation), 53, 234, 717–718
Morando, Mariana, Luciano J. Avila, and Jack W. Sites Jr., Sampling strategies for delimiting species: genes, individuals, and populations in the *Liolaemus elongatus–kriegi* complex (Squamata: Liolaemidae) in Andean–Patagonian South America, 159–185
Morphological
analysis, 187, 213, 323, 501, 504, 541, 560, 605, 858
characters, 55, 57, 70, 72, 74, 87–88, 89, 91, 131–132, 135, 142, 178, 181, 199, 209, 213, 217, 219, 298, 313–314, 317, 323, 327–328, 340, 389–390, 400, 406–407, 409, 412, 420, 439, 442, 457, 474, 501–503, 506, 509–510, 537, 539–543, 545, 559, 572, 604, 632, 738, 742, 748, 760, 855–857
data, 16, 20, 74, 103, 143, 182, 186–187, 243, 305, 315, 323, 329, 339, 369, 387–388, 397, 400–401, 403, 406, 423, 439, 501–503, 505, 726–727, 750, 764, 767, 771, 773, 796, 853–854, 857
evolution, 186–205, 404, 542
phylogeny, 206
studies, 727
support, 614–615
trees, 387, 392, 394, 396, 503, 860
Morphology, 10, 12, 16–17, 23, 57, 206–228, 239–252, 253–254, 296, 307, 318, 323, 328, 333, 477, 488, 501–514, 539–548, 559, 572, 614, 619, 632, 691, 748, 775, 851–852
genitalic, 71, 82
Morphometric
analysis, 197
data, 726
distance, 497
Morphometrics
geometric, 66–69, 186–205, 488–500
Morrison, David A., Branch lengths and support: revisited, 849–851
Moschidae, 206–228
Moschinae, 220
*Moschiola*, 206
*Moschus*, 206–207, 209, 212, 215, 219–221, 224
*moschiferus*, 210, 219
Mosses, 430
Most–parsimonius,
combined tree, 16
trees, 3, 5–7, 15, 55–60, 63, 575, 609, 621, 801, 804, 809, 826, 857–857, 864
Most recent common ancestor (MRCA), 48–49, 51–52, 179–180
Moths, tortricine, 430
*Mourasuchus atopus*, 417
Mouse, 290, 571, 591, 615
harvest, 677
Mouse deer, 206, 854, 857
Mouse lemur, 705–706, 708, 711–715
mtLSU-rRNA, 735–738, 740–741
mtSSU-rRNA, 735–738, 740
Multidimensional scaling (MDS), 190, 194, 693, 718–719, 721
Multigene analysis, 604
Multiple trees, 849–851
Multistate characters, 26, 75, 239, 255–257, 335
Multivariate analysis, 652
Muntiacinae, 209
*Muntiacus*, 209, 212, 215, 218–219, 222
*reevesi*, 210
Muridae, 606, 608
Muroidea, 604–605, 609, 613–614
Muroid phylogeny, 613
*Murrayona*, 314–315, 317, 324, 332–333
*axinellae*, 316
Murrayonidae, 314, 318, 329
Murrell, Anna, Nick J. H. Campbell, and Stephen C. Barker, Value of idiosyncratic markers and changes to conserved tRNA sequences from the mitochondrial genome of hard ticks (Acari: Ixodida: Ixodidae) for phylogenetic inference, 296–310
*Mus*
*domesticus*, 606
*musculus*, 286, 316, 736
*Musca*, 751, 753–754
*domestica*, 756
Muscidae, 756
Muscinae, 520
Muscomorpha, 746, 750–751, 753
Musk deer, 206–207, 210, 219–220
*Musophaga violacea*, 388
Mussels, 59
*Mustela*, 590
*erminea*, 573

*frenata*, 573, 577
*vison*, 573, 577
Mustelidae, 571–593, 846
Mustelinae, 571–573, 581
*Mycale fibrexillis*, 316
*Mydas* sp., 756
*Mydaus*, 573, 590
Mydidae, 750, 756
Myers, Daniel S., *see* Cummings, Michael P., 477–487
Myodonta, 604–606, 608, 610–611, 613, 615
Myomorpha, 614
Myriapoda, 557
*Mystacina*, 462, 464, 467, 473
*aupourica*, 460, 474
*gigas*, 473
*rhyacobia*, 460, 474
*tuberculata*, 460–463, 466, 470–474

*NADH*, 460
*Naja*, 447, 451
*kaouthia*, 441
*naja*, 441
National Science Foundation, 428–435
Natricinae, 439–440, 447, 452, 455–457
*Natrix*, 447, 451
*natrix*, 440
Naylor, Gavin J. P., and Dean C. Adams, Total evidence versus relative evidence: a response to O'Leary et al. (2003), 864–865
*ND1*, 73–74, 76, 657, 659
*ND2*, 460, 462–468, 471, 657, 659, 758–759, 762, 777
*ND3*, 657, 659
*ND4*, 161, 163–165, 167, 181, 440, 443, 445–447, 451, 453, 455, 657, 659
*ND4l*, 657, 659
*ND5*, 656–657, 659
*ND6*, 406, 657–659
Nearest Neighbor Interchange (NNI), 120, 698, 700
branch-swapping, 371
Near-optimal trees, 696
Neighbor joining (NJ), 121, 229–238, 368, 371–372, 506, 697, 699–702
analysis, 463, 530, 532
methods, 290, 293
tree, 211, 370, 372, 455, 462–463, 508, 702, 801
Nematocera, 754
Nematodes, 430
Nemerteans, 430
Nemestrinidae, 746, 753–754
Nemestrinids, 753
*Neoaetosauroides*, 241, 243, 248, 250–251
*Neoceratina*, 2
Neogene, 206
Neologisms, 515, 517
*Neorhycocephalus volaticus*, 756
*Neotragus pygmaeus*, 221
*Neotrigonia*, 559
*Nephila plumipes*, 300
Nepokroeff, Molly, Kenneth J. Sytsma, Warren L. Wagner, and Elizabeth A. Zimmer, Reconstructing ancestral patterns of colonization and dispersal in the Hawaiian understory tree genus Psychotria (Rubiaceae): a comparison of parsimony and likelihood approaches, 820–841
*Neriene*, 72, 77, 82
*radiata*, 73–74, 77, 88
*variabilis*, 73–74, 77, 88
*Nerita*, 317
*albicilla*, 316
Nerodia, 447, 451
*fasciata*, 440
*rhombifera*, 440
*taxispilota*, 440
*Nesobasis*, 91, 94, 96
*Nesomantis thomasseti*, 287
Nested-clade analysis (NCA), 159–185
Nested model, 606
Newton–Raphson
algorithm, 699
method, 698, 703
New World, 207, 548–549, 552
*Crocodylus*, 409
ground doves, 39
lizards, 160
New Zealand pigeon, 45
*Nicotiana*, 347
Nielsen, Rasmus, *see* Huelsenbeck, John P., 131–158
*Noctilio*, 471
*albiventris*, 464
*leporinus*, 464
spp., 463
Noctilionoidea, 460, 464
Nonhomogeneous maximum likelihood analysis, 390
Nonparametric
bootstrap, 506, 594, 665–667, 669, 671–672, 726, 760, 790, 801
bootstrap analysis, 165
rate smoothing method (NPRS), 76
sign-rank test, 16
test, 781
Nontrivial tree, 485
Norell, Mark, *see* Gatesy, John, 403–422
*Nosomma*, 305
*monstrosum*, 299, 305
*Nothofagus*, 160
Novacek, Michael J., see O'Leary, Maureen A., 853–866
Novel versus unsupported clades: assessing the qualitative support for clades in MRP supertrees, Olaf R. P. Bininda-Emonds, 839–851
*Novosieversia glacialis*, 382
nrDNA, 336
*Nucella*, 631
Nuclear
alleles, 41, 103
DNA, 37, 89, 95, 99, 103, 221, 374–375, 382–383, 407–408, 415, 418, 423, 509, 571–593, 735, 757–758, 760–762, 764, 767, 770–771, 773, 775–776
genes, 16, 28–29, 32, 41, 74, 375, 386, 572
genome, 374–375, 571, 604
sequence, 308
substitution model, 215, 287, 377, 604–617, 658, 674–683, 708
Nucleotides, 122, 124, 126, 296, 300–301, 303, 692
data, 305
sequence, 156, 697, 747
substitution, 121, 135, 375, 445, 465, 478, 480, 621, 651, 655, 750
substitution model, 377, 604–617, 658
Null
distribution, 30, 761
hypothesis, 16, 28, 241, 249, 278–279, 363–364, 425, 464, 478, 480, 598, 600, 607, 609, 611, 613, 718, 720, 751, 761, 770, 775, 781–784, 788, 790–792, 801
model, 271–280, 480, 482, 484, 717–723
Nyctaginaceae, 334, 344, 346–348
Nyffeler, Reto, *see* Mast, Austin R., 271–280
Nymphalid butterflies, 28

Occam's window, 50
*Occidozyga alevis*, 719, 815
*Ochotona*, 609
*princeps*, 606
Odocoileini, 209
*Odocoileus*, 209, 212, 215, 218–219
*hemionus*, 210
*virginianus*, 210
Odonata, 89
*Oena capensis*, 38, 46
Okapi, 206, 210
*Okapia*, 207, 209, 212, 218–219, 221–222, 224
*johnstoni*, 210
Old World, 206–207, 548–550, 552
Arenaviridae, 740
badgers, 577, 590
O'Leary, Maureen A., John Gatesy, and Michael J. Novacek, Are the dental data really at odds with the molecular data? Morphological evidence for the whale phylogeny (re)reexamined, 853–866
Oligocene, 224–225, 795
early-, 206–207, 216–218, 224
late-, 206–207, 216–218, 225
Oligocephalus, 492
Olmstead, Richard G., *see* Scotland, Robert W., 539–547
Onagraceae, 347
*Oncostylus*, 382
*cocklaynei*, 381
*leiospermus*, 381
One-tailed test, 28, 339
Ontogeny, 430, 515–527
Onycophora, 557
*Oopsacas minuta*, 316
Operational taxonomic units (OTU), 291, 620
*Ophidiocephalus*, 757–758, 764, 767, 770–771
taeniatus, 757, 759
*Ophiophagus*, 451
*hannah*, 441
Opilioacarid, 300, 309
Opilioacarida, 300
Opilioacariformes, 300
Opiliones, 557
Opossum, 285, 656
Opsin, 28–30
Optimal tree, 4, 8, 17, 20, 121, 128, 320, 341, 357, 696, 760, 850
Orangutan, 52–53, 794
Orchidaceae, 346
*Oreamnos*, 225
*Oreocalamus*, 447, 451, 457
*hanitschi*, 440
*Oreotragus*, 221
Ornithodorinae, 300
*Ornithorhynchus anatinus*, 285
*Oropsylla silantiewi*, 187
*Orsonwelles*, 70–88, 103, 833
*ambersonorum*, 73, 84, 88
*arcanus*, 73, 79, 84, 88
*bellum*, 71, 73, 88
*calx*, 73, 84, 88
*falstaffius*, 73, 79, 84, 88
*graphicus*, 71, 73, 79, 84, 88
*indicium*, 71, 73, 79, 88
*macbeth*, 73, 77, 79, 84, 88
*malus*, 71–73, 77, 79, 81, 88
*othello*, 73, 77, 79, 84, 88
*polites*, 72–73, 79, 84, 88
*torosus*, 71, 76–77, 79, 81–82, 88
*ventus*, 73, 79, 88
*Ortalis*,
*cinereiceps*, 388
*guttata*, 286
Orthoptera, 307
*Orthotylus*, 103
*Oryctolagus cuniculus*, 286, 289, 606
Osteoderms, 239–252, 253–255, 851–852
*Osteolaemus*, 386, 391, 409
*tetraspis*, 388, 391, 405, 408–409, 415
Osteological characters, 760
Ostrich, 388
Otariidae, 846
*Otobius megnini*, 300
*Otospermophilus*, 197
*variegatus*, 189
Otters, 571–573, 590
Asian small-clawed, 573
Cape clawless, 573
Eurasian, 573
giant, 573
marine, 573
Neotropical, 573
North American, 573
river, 577, 590
sea, 573
spotted-necked, 573
*Ourebia ourebi*, 221
Outgroup, 9, 27, 39, 48, 52, 56, 60–61, 81–82, 88, 89, 92, 94, 96, 99, 102, 189, 192, 196, 205, 229–230, 232, 235, 237, 298, 301, 309, 386–387, 391–392, 404, 415, 443, 446, 463–464, 557, 574, 576, 618, 620–622, 660–661, 674, 707, 756–757, 762, 764, 767, 770–771, 775, 798–799, 815, 821, 826–827
Outgroup misplacement and phylogenetic inaccuracy under a molecular clock—a simulation study, B. R. Holland, D. Penny, and M. D. Hendy, 229–238
*Ovibos*, 225
*Ovis*, 209, 215, 218, 225
*aries*, 210, 212
Owl monkeys, 738
Oxelman, Bengt, *see* Erixon, Per, 665–674
*Oxylipeurus chiniri*, 39

*Pachygaster leachii*, 756
Paedomorphic salamanders, 503
Page, Roderic D. M., *see* Johnson, Kevin P., 37–47
Pairwise tests, 339, 344
*Palaeobolbomyia*, 746, 750
Palaeoheterodonta, 559–560
*Palaeoplatypygus*, 749
Paleocene, 551
late-, 550
Paleontological analysis, 605
Paleontology, 206–207, 265
*Paleosuchus*, 391, 409
*palpebrosus*, 388, 405
*trigonatus*, 405
*Pan*, 706
Pantophthalmidae, 756
*Pantophthalmus* sp., 756
*Papilio*, 589
*Paradelma*, 758, 775
*orientalis*, 758, 761
Paradialictus, 24, 26
*Paralamyctes*, 559

Paralictus, 24
Paralinear distances (*see also* LogDet/paralinear distance), 287, 291
*Paraleucilla*, 314–315, 320, 322, 324–325, 332
  sp., 316
Parallel evolution, 296, 307
Parametric
  bootstrap, 18, 598, 667, 726, 757–780
  bootstrap test, 607, 802, 807, 809, 812
  test, 790
*Paraplatypeza atra*, 756
Parasites, 26, 37–43, 45–46, 406, 735, 737–738, 740–742
  phylogeny, 740–741
  tree, 740, 742
Parasitiformes, 299
Parasitoids wasps, 430
Parasphecodes, 24
*Paratomistoma*, 399
*Paratypothorax*, 241, 243, 248–252
Paraxonia, 853, 861–862
*Parazoanthus*, 317
  *axineliae*, 316
*Pareas*, 451
  *nuchalis*, 440, 446
Pareatinae, 439–440, 447, 452, 455
Parisitological characters, 738, 742
Parrots, 549–553
Parsimony (*see also* maximum parsimony), 19, 23, 39, 41, 76, 122, 124, 126, 131–132, 140, 147, 236, 258, 265, 335, 362–367, 386, 394, 396, 501, 556, 584, 604, 632, 641–648, 656, 696, 750, 761, 809, 820, 827, 832, 834–835, 841
  algorithms, 867
  analysis, 7, 26, 30, 42, 45, 70, 76, 79, 83, 249, 251, 255, 257, 260, 262, 271–280, 336, 377, 389–390, 392, 406, 412, 423, 506, 508, 530, 554, 571, 575, 583, 618, 620–621, 623, 629, 727, 764, 802, 804, 824
  analysis of endemicity (PAE), 271–280
  Camin–Sokal, 444, 446
  cladistic, 356, 365
  equally weighted, 620–621
  Fitch, 444, 446
  informative characters, 4, 323, 345, 446, 451, 529, 531, 773
  informative indels, 823
  informative sites, 28, 30, 613
  jackknife, 128
  jackknife support, 59, 557, 559
  linear, 342
  matrix, 241
  matrix-weighted, 623
  model, 361, 363
  permutation tail probability (PTP), 250, 252
  ratchet 368–373
  reconstruction, 629
  search, 368, 842
  squared-change, 337, 342
  support, 623
  trees, 9–10, 39, 42–43, 79, 286, 645, 762, 764, 771, 773, 802–804
  uniformative characters, 298
  weighted, 284
Parsons, Thomas J., *see* Harshman, John, 386–402
Partition analysis, 4
Partitioned
  analysis, 396, 604–607
  branch support (PBS), 15–22, 76, 79, 307, 407, 409, 412
  likelihood analysis, 386–402
  likelihood support (PLS), 15–22,
Partitioned likelihood support and the evaluation of data set conflict, M. S. Y. Lee and A. F. Hugall, 15–22
Partition homogeneity test (PHT; *see also* incongruence length difference [ILD] test), 6, 15, 41–42, 45, 344, 347
Passeriformes, 552
*Patagioenas*
  *fasciata*, 38
  *plumbea*, 38
  *speciosa*, 38
  *subvinacea*, 38
PcP, 735–744
*p*-distances, 5
Pecora, 206–207, 209, 213, 218, 221–222, 224–225
*Pedetes*, 613
  *capensis*, 606
Pedetidae, 604–606, 610–611, 613–614
*Pelagomapouria*, 826, 829, 832–834
Pelecorhynchidae, 753, 756
*Pelecorhynchus personatus*, 756
Penny, D., *see* Holland, B. R., 229–238
Peracarida, 430
Percidae, 488
*Percina*, 491
  *caprodes*, 497, 500
  *nigrofasciata*, 491, 497, 500
  *ouachitae*, 497, 500
  *roanoka*, 491, 497–498, 500
  *sciera*, 491, 497, 500
  s.l., 494, 497
Performance-based selection of likelihood models for phylogeny estimation, Vladimir Minin, Zaid Abdo, Paul Joyce, and Jack Sullivan, 674–683
*Pericharax*, 314–315, 332
Periods,
  cenozoic, 206, 631, 746
  cretaceous, 23, 33, 403, 419, 550–552, 631, 746, 753, 861
  early Cretaceous, 548–549
  early Eocene, 795
  early Miocene, 206–207, 209, 217–218, 224–225
  early Oligocene, 206–207, 216–217, 224–225
  eocene, 222, 403, 419, 857–858
  jurassic, 746, 751, 753
  late Cretaceous, 419, 549–552
  late Eocene, 222, 224
  late Jurassic, 549, 746
  late Miocene, 218, 224
  late Oligocene, 225
  late Paleocene, 550
  late Pleistocene, 473
  late Triassic, 549, 745, 751, 754
  lower Cretaceous, 753
  lower Jurassic, 746, 749
  lower/middle Triassic, 746
  mesozoic, 549–553, 745–746
  mid-Cretaceous, 549, 552
  mid-Eocene, 218, 222, 224–225
  mid-Jurassic, 745
  mid-Miocene, 206–207, 209, 217–218, 224–225
  mid-Pleistocene, 198, 460, 473, 810
  mid-Tertiary, 549
  mid-Triassic, 548
  miocene, 207, 221, 451, 549, 551, 629, 761, 776, 808
  neogene, 206
  oligocene, 224–225, 795
  paleocene, 551
  pleistocene, 186, 198, 460–476, 777, 797, 807–808, 812
  pliocene, 224–225, 629
  plio-pleistocene, 225

quaternary, 552
tertiary, 549–553, 746, 753
triassic, 239–252, 403, 749, 751
upper Cretaceous, 753
upper Jurassic, 753
upper Triassic, 753
Perissodactyla, 850, 857–858
Permutation tail probabilities (PTPs), 250, 252
*Petrobiona*, 314–315, 317, 320, 322–324, 328–329, 332–333
*massiliana*, 316
Petrobionidae, 314
*Petromarmota*, 186, 192–195, 197–198
*caligata*, 189
*flaviventris*, 189
*olympus*, 189, 193
*vancouverensis*, 189
*Petrostoma*, 329
*Phapitreron*
*amethystina*, 38
*leucotis*, 38
Phenotypes, 241, 349
Philippines, 794–819
Philosophy, 352–367
of science, 259–271, 352
Philothamnini, 457
Phocidae, 846–847
Phthiraptera, 37–47
Phyllostomoidea, 460
Phylogenesis, 520
Phylogenetically informative characters, 55–56, 826
Phylogenetic
accuracy, 121, 528–538
artifacts, 594
clade, 714
comparative method (PCM), 684–695
congruence, 165
conservatism, 1
data, 89, 248–249, 314, 649, 724, 726, 745, 747
distance, 491
effect, 488–500, 684, 686, 692, 694
eigenvector regression (PVR), 694
error, 119, 121, 124–125
estimates, 159, 284, 530, 802, 850
evidence, 503
framework, 311, 728
generalized least squares (PGLS),
history, 91, 131–132, 337, 348–349, 501, 589–590, 641, 745
hypothesis, 1, 55, 62, 212, 268, 283, 352, 354, 362–363, 401–406, 445, 477, 488, 554, 556–557, 571, 632, 672, 689, 693, 725–726, 728, 740, 742, 745, 764, 854, 858, 860–861
inference, 75, 180, 187, 296, 306, 335, 344, 352, 359, 368–373, 463, 511, 554, 612, 641, 696, 791
information, 96, 296, 311, 327, 339, 589, 686, 747, 761
level, 542
markers, 864
methods, 131, 229, 283, 285, 291, 293, 352, 358, 469, 480, 641–642, 665, 677
model, 133, 594, 649, 651, 654, 656, 661
patterns, 342–344, 348
reconstruction, 19, 39, 83, 287, 318, 375, 539–548, 571, 658, 665, 669, 671, 685, 724–729, 860–861, 864
relationships, 2, 23, 35, 71, 92, 102, 159, 179, 181, 186, 199, 213, 215, 220, 239, 245, 255, 265, 335–336, 347, 439, 444, 447, 451, 471, 477, 548, 554, 591, 623, 728, 745, 775, 797, 828
signal, 39, 55, 63, 70, 79, 83, 186, 197, 199, 328, 334, 371–372, 403, 412, 415–416, 445, 497, 511, 577, 581, 584, 591, 773, 853–855, 857, 860
species, 833
structure, 178, 181, 473, 686–687, 691
studies, 32, 110, 188, 374–375, 545, 757–758, 821–822
supertrees, 839–851
systematics, 260, 510, 735
taxonomy, 852
tests, 781
tools, 311
tree, 119, 121, 126, 71, 112, 131, 134–135, 244, 289–290, 359, 364–365, 427, 463, 515, 554–555, 669, 671, 685, 692, 715, 749, 760–761, 771, 791, 824, 847, 849
uncertainty, 39, 132, 488
utility, 27, 30, 32
Phylogenetic methods and Aetosaur interrelationships: a rejoinder, Simon R. Harris, David J. Gower, and Mark Wilkinson, 851–852
Phylogenetic relationships among Calyptraeid gastropods and their implications for the biogeography of marine speciation, Rachel Collin, 618–640
Phylogenetic systematics and evolution of primate–derived *Pneumocystis* based on mitochondrial or nuclear DNA sequence comparison, Jean–Pierre Hugot, Christine Demanche, Véronique Barriel, Eduardo Dei–Cas, and Jacques Guillot, 735–744
Phylogenetics, 1–14, 48, 70–88, 95, 127, 136, 243, 247, 270, 334–351, 352–367, 386, 477–487, 515–527, 557, 565, 594–603, 655, 665, 782–783, 794, 847, 862
Bayesian, 48, 54, 89–109
molecular, 187, 283–295, 311, 715
vertebrate, 388
Phylogenetics of advanced snakes (Caenophidia) based on four mitochondrial genes, Christopher M. R. Kelly, Nigel P. Barker, and Martin H. Villet, 439–459
Phylogenetics of fanged frogs: testing biogeographical hypotheses at the interface of the Asian and Australian faunal zones, Ben J. Evans, Rafe M. Brown, Jimmy A. McGuire, Jatna Supriatna, Noviar Andayani, Arvin Diesmos, Djoko Iskandar, Don J. Melnick, and David C. Cannatella, 794–823
Phylogenomics, 119–124
Phylogeny, 8–9, 23–36, 37–47, 48–54, 57, 60–61, 63, 66, 89, 92, 111, 114, 121, 133, 136, 139, 152, 159–185, 186–205, 206–228, 229, 239, 253–255, 264, 272, 280, 296–310, 311–333, 334–335, 345, 348, 368, 374–385, 386–402, 430, 460–461, 471, 488, 493, 497–498, 501–514, 515, 520–521, 525–526, 539, 543, 548, 571–593, 604–617, 632, 645, 647, 649–664, 665–673, 674–683, 687, 692–693, 696–704, 714, 724, 726, 728, 735, 737, 740, 751, 757–780, 792, 798, 801–802, 820, 855, 864
aetosaur, 852
cetacean, 854
chloroplast, 375
estimates, 423–428, 674, 677, 682
fly, 746
gene, 374
host, 740, 742
hypotheses, 738
mammalian, 857
molecular, 37–47, 206, 545, 781, 832
morphological, 206
muroid, 613
parasite, 740–741
*Pneumocystis*, 737, 740
reconstruction, 264, 349, 501, 511, 542–543, 545, 586
rodent, 613
tetrapod, 283–295
whale, 853
Phylogeny and evolution of Calcareous sponges: monophyly of Calcinea and Calcaronea, high level of morphological

homoplasy, and the primitive nature of axial symmetry, Michael Manuel, Carole Borchiellini, Eliane Alivon, Yannick Le Parco, Jean Vacelet, and Nicole Boury-Esnault, 311–333
Phylogeny of eusocial *Lasioglossum* reveals multiple losses of eusociality within a primitively eusocial clade of bees (Hymenoptera: Halictidae), Bryan N. Danforth, Lindsay Conway, and Shuqing Ji, 23–36
Phylogeny reconstruction: the role of morphology, Robert W. Scotland, Richard G. Olmstead, and Jonathan R. Bennett, 539–547
Phylogeographic
analysis, 460
structure, 460–476
Phylogeography, 92, 801, 812
Phylogram, 100, 140, 621
*Phymaturus indistinctus*, 160, 164
*Physeter*, 706, 712
Phytosaurs, 253
Pianka, Eric R., *see* Jennings, W. Bryan, 757–780
*Pichia anomala*, 316
*Picoides pubescens*, 286
Pigeons, 37, 41, 45
New Zealand, 45
Pigs, 854
Pinworm, 742
*Pithecia*, 737, 740, 742
*pithecia*, 736
*Pityohyphantes*, 77
*costatus*, 73, 88
Placental mammals, 591
Placozoa, 31
*Plagithmysus*, 89
Plankton, 618
Plants, 89, 106, 335, 347, 517, 541, 696, 746, 794, 820
flowering, 375, 518, 753, 820
seed, 542, 781
Plasmid DNA, 316
Platodes, 520
Platymantinae, 794
Platypezidae, 756
Platypodid beetles, 23
Platyrrhini, 738, 742, 735, 737, 741
Plecoptera, 106
*Plectroninia*, 314–315, 317, 323, 328–329, 332
*halli*, 329
Pleistocene, 186, 198, 777, 797, 807–808, 812
divergence, 460–476
late-, 473
mid-, 198, 460, 473, 810
Plesiometacarpalia, 207, 215
Plethodontid salamanders, 510
*Pletholax*
*gracilis*, 757, 759
*hortulanus*, 761
Pliocene, 629, 629
Plio-pleistocene, 225
*Pneumocystis*, 735–744
*carinii*, 735–736
*carinii aotus*, 736, 738, 742
*carinii callimico*, 736, 738
*carinii callithrix*, 736, 738, 740
*carinii fascicularis*, 736
*carinii fuscicollis*, 736, 738
*carinii geoffroyi*, 736, 738, 740
*carinii griseus*, 736
*carinii hamlyni*, 736
*carinii imperator*, 736, 738
*carinii macaco*, 736
*carinii midas*, 736, 738
*carinii mulatta 1*, 736
*carinii mulatta 2*, 736
*carinii muris*, 736
*carinii nemestrina*, 736
*carinii nictitans*, 736
*carinii nigroviridis*, 736
*carinii oedipus*, 736
*carinii pithecia*, 736, 738, 742
*carinii sciureus*, 736, 738, 742
DNA, 736
genes, 742
genotypes, 735
*jiroveci*, 736
phylogeny, 737, 740
pneumonia (PcP), 735–744
spp., 735, 738, 742
Poe, Steven, Evolution of the strategy of long-branch subdivision to improve the accuracy of phylogenetic methods, 423–427
Poe, Steven, *see* de Queiroz, Kevin, 352–367
Poisson
distribution, 133
process model, 134
Polhemus, Dan, *see* Jordan, Steve, 89–109
*Polinices didyma*, 640
Pollock, David D., *see* Hillis, David M., 124–126
Polychaetes, 430
Polymerase chain reaction (PCR), 26, 39, 74, 94, 301, 316, 376, 383, 389, 404, 571, 574, 591, 606, 620, 736, 738, 758, 823
amplification, 2–3, 161, 375, 389, 736, 747
recombination, 384
Polymorphic characters, 505, 833
*Pongo*, 706
*Pontederia*
*cordata* var. *cordata*, 154, 156
*cordata* var. *lancifola*, 154, 156
*cordata* var. *ovalis*, 154, 156
*rotundifolia*, 154, 156
*sagittata*, 154, 156
Pontederiaceae, 139–140, 143, 147, 154, 156
Popper and systematics, Olivier Rieppel, 259–270
Popper, Karl, 259–271, 352–367
Popper's degree of corroboration, 352, 355, 360, 363, 366
Porifera, 311–333
Potent antiretroviral therapy (PART), 660–661
Prairie dog, 186,
Prestosuchidae, 252
*Priapulus caudatus*, 316
Price, Samantha E., *see* Bininda-Emonds, Olaf R. P., 724–729
Primary characters, 725–726
Primates, 52–53, 57, 549, 606, 707–708, 735–744
Principle component analysis (PCA), 687
Principle coordinates analysis, 860
*Pristichampsus*, 409
Probability, 352–367
*Procervulus*, 224
Procrustes, 66–69
distance, 67, 190, 192, 493
generalized analysis, 66, 68
geometry, 67–68
Procyonidae, 573–574, 846
Procyoninae, 573
*Procyon lotor*, 573–574

Progressive rearing, 1–14
Prokaryotes, 658–660
*Propithecus*, 706
Prosobranchia, 328
Prostriata, 296, 300, 304, 307
Protein-coding, 391
  genes, 23, 39, 74, 94, 297–298, 307, 358, 396, 405, 445, 460, 471, 506, 604, 609, 612, 656, 682, 705–706
  regions, 52, 165
Protein evolution, 602
Protists, 429–430
*Protragocerus labidotus*, 221
Prunoideae, 378–379, 382
*psaA*, 785, 787, 791
*psaB*, 785, 787–788, 791
Psammophiinae, 439–440, 452, 455
*Psammophis*, 451
  *condenarus*, 441
  *mossambicus*, 441, 459
*Psammophylax*, 451
  *variabilis*, 441, 459
*Pseudagrion*, 91, 94, 102
  *civicum*, 94
  *pacificum*, 94
  *palauense*, 94
*Pseudemys scripta*, 286–287
*Pseudochilalictus*, 24
Pseudoboini, 456
Pseudoreplicate, 55–65
Pseudoxyrophiinae, 439, 441, 443, 452, 455–456
Psittacidae, 551
Psittiformes, 551
Psychodomorpha, 756
*Psychomyia*,
*Psychotria*, 820–841
  *cadigiensis*, 821–822
  *fauriei*, 822, 826, 829, 833–834
  *grandiflora*, 822, 826, 828–829, 832, 834
  *greenwelliae*, 820, 822–823, 825–827, 829, 832–834
  *hathewayi*, 822, 826, 829, 833–834
  *hawaiiensis* subsp. *hawaiiensis*, 829
  *hawaiiensis* var. *hawaiiensis*, 822, 829, 833
  *hawaiiensis* var. *hillebrandii*, 822–823, 829
  *hexandra*, 822, 828–829, 832, 834
  *hexandra* subsp. *O'ahuensis*, 822
  *hobdyi*, 822, 828–829, 832, 834
  *hombroniana*, 821–822
  *kaduana*, 822, 829, 831, 833
  *luzionensis*, 821–822
  *mariniana*, 822, 825, 827, 829, 832–834
  *mauiensis*, 822, 829, 833
  *pickeringii*, 821–822
  sp. Fiji, 821–822
  sp. Guam, 822
  sp. Tinian Is., 821
  *tahitiensis*, 821–822, 826
  *wawrae*, 822, 829, 834
Pteriomorphia, 561
*Pteronura*, 590
  *brasiliensis*, 581, 583, 589–590
Pterygota, 328
*Ptilinopus occipitalis*, 38
*Ptiolina fasciata*, 756
Punctuational evolution, 831
Pupal characters, 8
Purvis, Andy, *see* Bininda-Emonds, Olaf R. P., 724–729
*P*-values, 18, 22, 28, 99, 132, 137, 147, 156–157, 190, 192, 273, 279, 345, 394–395, 406, 425, 598–600, 692, 722, 761, 770, 783–785, 787–788, 790–791
  adjustment, 781–793
Pycnogonida, 557
Pygopodidae, 757–780
Pygopodid
  lizards, 757–780
  phylogeny, 758, 764
  tree, 757, 760, 770
*Pygopus*, 757–758, 762, 764, 767, 770–771, 775
  *nigriceps*, 759–761, 775
  (*Paradelma*) *orientalis*, 759
  *schraderi*, 758
  *steelescotti*, 758
Pythonidae, 440
*Python reticulatus*, 440

Quail, 551
Qualitative support (*QS*) index, 839, 841, 844, 847
*Quamoclidion*, 336
Quartet puzzling (QP) values, 558
Quaternary, 552

Rabbits, 289
Raccoon, 573
RAG-1, 405, 408
RAG-2, 604, 606–610, 613–614
Raguso, Robert A., *see* Levin, Rachel A., 334–351
*Rana*
  *nigromaculata*, 798
  *signata*, 810
  *vittigera*, 815
Random
  generation, 110–118
  trees, 30, 115, 445, 681
*Rangifer*, 209, 212, 215, 218–219
  *tarandus*, 210
Rangiferini, 209
*Ranodon sibiricus*, 286
Ranunculaceae, 347
*Ranunculus*, 347
Rare genomic changes, 296
Rate
  distribution, 594–603
  homogeneity model, 463
  of evolution, 602, 641, 646–647, 677, 746, 753, 833
Rates-across-sites model, 595–596
Ratite, 290
  birds, 549–553
Rats, 290
*Rattus norvegicus*, 286, 606, 736
Rauisuchia, 241–243, 251
*rbc*L, 140, 336, 379, 696–704
rDNA, 405, 408, 808, 820
Recapitulation, 515–527
Recognition, 110–118
Reconciliation analysis, 39, 42
Recursion, 110–118
Reconstructing ancestral patterns of colonization and dispersal in the Hawaiian understory tree genus Psychotria (Rubiaceae): a comparison of parsimony and likelihood approaches, Molly Nepokroeff, Kenneth J. Sytsma, Warren L. Wagner, and Elizabeth A. Zimmer, 820–841
RDPII project, 696–704
Red deer, 210

*Redondasuchus*, 241–243, 248–252
Red panda, 846
Reductive characters, 245
*Redunca*, 209, 212, 215
  *fulvorufula*, 210
Reduncini, 209–210
Redundancy analysis (RDA), 687–689, 693
Reed, David L., *see* Cummings, Michael P., 477–487
Regression, 66–69
  analysis, 68
  model, 684, 686
Regularia, 328
Reindeer, 210
*Reithrodontomys sumichrasti*, 677
Relative warp analysis (RWA), 198
Reliability of Bayesian posterior possibilities and bootstrap frequencies in phylogenetics, Per Erixon, Bodil Svennblad, Tom Britton, and Bengt Oxelman, 665–673
RELL bootstrapping, 390
Remipedia, 557
Reptiles, 524
Reptilia, 239, 253
Rescaled consistency index, 762, 764
Retention index, 3, 5–6, 10, 28, 56, 76, 217, 304, 339, 345–346, 348, 577–578, 581–583, 762, 764, 826
Reticulate evolution, 374–385
Reweighted characters, 318, 327, 371
Rhagionidae, 746, 756
*Rhagoletis*, 754
Rhamnaceae, 375
*Rhamnus*, 378–379
  *cathartica*, 377
Rhamphiophis, 451
  *acutus*, 441, 459
  *oxyrhynchus*, 441
*Rhamphotyphlops braminus*, 440
*Rhea americana*, 550
Rheas, 550
*Rhingia nasica*, 756
Rhinocerus, 706–708, 794
*Rhipicentor*, 305–306
  *nuttalli*, 299
Rhipicephalinae, 297–298, 305–307
  s.l., 296, 299, 304–305, 307, 309
*Rhipicephalus*, 299, 306
  *appendiculatus*, 299, 306, 310
  *compositus*, 299
  *evertsi*, 299, 310
  *maculatus*, 299
  *pravus*, 299, 310
  *pulchellus*, 299, 306, 310
  *pumilio*, 306
  *punctatus*, 299
  *sanguineus*, 299, 306, 310
  *simus*, 299
  *turanicus*, 299, 306
  *zambeziensis*, 299
  *zumpti*, 299
*Rhizomys*, 613
  *pruinosus*, 606
Rhynchosaurs, 253
Rieppel, Olivier, Popper and systematics, 259–270
Ringtail, 573
Rivulidae, 548, 552
*Rivulus marmaratus*, 549
RNA, 555, 610
  mitochondrial, 445
  polymerase II, 557
  secondary structure, 283–295
  sequencing, 49, 651
Rock squirrel, 194, 197
Rodent, 656, 679–680
  Holarctic, 186
  phylogeny, 613
  tree, 615
Rodentia, 186, 604–617
Rodman, James E. and Jeannine H. Cody, Taxonomic impediment overcome: NSF's Partnerships for Enhancing Expertise in Taxonomy (PEET) as a model, 428–435
Roger, Andrew J., *see* Susko, Edward, 594–603
Rohlf, F. James and Fred Bookstein, Computing the uniform component of shape variation, 66–69
Rokas, Antonis, *see* Cummings, Michael P., 477–487
Rooted
  duplication tree (RDT), 110–111, 113, 115
  tree, 865
Rooting, 386–402
Roots of phylogeny: how did Haeckel build his trees?, Benoît Dayrat, 515–527
*Rosa*, 375
  *multiflora*, 375–377
Rosaceae, 374–377, 379, 382
Rosenberg, Michael S. and Sudhir Kumar, Taxon sampling, bioinformatics, and phylogenomics, 119–123
Rosoideae, 378–379
  rRNA, 131, 283–285, 291, 293, 297, 307, 615
Rubiaceae, 820
*Rubus*, 375
  *odoratus*, 376–377
Ruminantia, 206–228
*Rupicapra rupicapra*, 221
Rutaceae, 347

Sable antelope, 210
*Saccharomyces cerevisiae*, 316
*Saguinus*, 740, 742
  *fuscicollis*, 736–737
  *imperator*, 736
  *midas*, 736, 738
  *oedipus*, 736
*Saimiri*, 740, 742
  *sciureus*, 736
Salamanders, 287, 501–514
  cave, 501
  hemidactyliine, 509
  paedomorphic, 503
  plethodontid, 510
Sampling design, 159–185
Sampling strategies for delimiting species: genes, individuals, and populations in the *Liolaemus elongatus–kriegi* complex (Squamata: Liolaemidae) in Andean–Patagonian South America, Mariana Morando, Luciano J. Avila, and Jack W. Sites Jr., 159–185
Sandvik, Hanno, Genes, categories, and species: the evolutionary and cognitive causes of the species problem (rev.), 565–567
Sarcoscyphineae, 430
*Sarona*, 833
Savage–Dickey ration, 48, 50, 53
Scale, 271–280
  of being, 515–527
*Sceloporus jarrovii*, 160
Scenopinidae, 750, 756

Scent, 334–351
*Schiedea*, 820, 832
*Schistosoma*, 307
Schizophora, 746, 749–751, 753–754
Schwarz, Michael P., Nicholas J. Bull, and Steven J. B. Cooper, Molecular phylogenetics of Allodapine bees, with implications for the evolution of sociality and progressive rearing, 1–14
Scincidae, 757
Sciuridae, 186, 605–606, 608, 614–615
Sciurid convergence, 186–205
Scuirids, 188, 197
Sciurini, 189
Sciuroidea, 604–606, 608–610, 613, 615
Sciuromorpha, 605, 614
*Sciurus*
  *infuscatus*, 205
  *vulgaris*, 189, 205
*Scobius*, 271–280, 717–723
  *aciculatifrons*, 273
  *cognatus*, 273
  *griseus*, 273, 275
  *horni*, 275
  *obesus*, 275
  *oneill*, 275
  *pullus*, 275
  *viridis*, 275
Scolecophidia, 440
*Scolopendra cingulata*, 316
Scorpiones, 300, 556–557
Scorpionoidea, 556
Scotland, Robert W., Richard G. Olmstead, and Jonathan R. Bennett, Phylogeny reconstruction: the role of morphology, 539–547
*Scypha*, 325
  *ciliata*, 317
Sea anemones, 430
Seastar, 139–140, 143
Sea urchins
  *Echinometra*, 631
Seed plants, 542, 781
Selenodontia, 221
*Sellalictus*, 24, 26, 30
*Semicassis pyrum*, 640
Sensitivity analysis, 554–564
Sequence analysis, 737
Sequence-tagged sites, 571–593
*Serinus canaria*, 389
Sex ratio, 24, 33
Shape statistics, 66–69
Sheep, 206, 210, 854, 857
Shell characters, 619, 623
Shimodaira–Hasegawa test,15–22, 96, 98, 212, 216, 221, 444–445, 464, 468, 504, 508, 781–793
Short-tailed bat, 460
Shrimp, 23
*Synalpheus*, 631
*Sicista*, 614
  *tianshanica*, 606
*Sieversia*, 374
*Sigapatella*, 619, 632, 639
  *novaezelandiae*, 639
  *terraenovae*, 639
Significance test, 781–793
Silicispongia, 320
Silverswords, 89
*Simimeryx*, 222
Simon, Chris, *see* Jordan, Steve, 89–109
Simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood, Stéphane Guindon and Olivier Gascuel, 696–704
Simulation (*see also* computer simulation and Monte Carlo simulation), 423–428, 665–673
Single-copy genes, 375
Single-gene tree, 20
Single-valve decomposition (SVD), 68–69
Sinsheimer, Janet S., *see* Suchard, Marc A., 48–54, 649–664
*Siphopatella*, 619, 623, 639
  *walshi*, 623, 639
Sites-specific rates (SSR)
  analysis, 612
  model, 4, 8, 26–28, 30, 463, 466–468, 610, 612, 674
  tree, 610–611
Sites, Jack W., Jr., *see* Morando, Mariana, 159–185
Skeletal characters, 199, 222, 313, 403
Skunks, 572–573
Smedmark, Jenny E. E., Torsten Eriksson, Rodger C. Evans, and Christopher S. Campbell, Ancient Allopolyploid speciation in Geinae (Rosaceae): evidence from nuclear granule-bound starch synthase (GBSSI) gene sequences, 374–385
Snails, 70
Snakes, 286–287, 456
  African, 456
  advanced, 439–459
  colubrid, 758
  Madagascan, 456
Snipe flies, 745–746
Snout characters, 416
Social
  behavior, 23–36
  evolution, 1–14, 23–36
Soft ticks, 300
Solanaceae, 347
Soldier flies, 746
Soleneiscidae, 314
*Soleneiscus*, 314–315, 317, 322, 329, 332–333
  *radovani*, 316
Sordariales, 430
Source trees, 727, 840–844, 846
South Africa, 717–723
Southern cassowary, 388
*Spalax*, 613
  *zemni*, 606
Spearman's rank correlation test, 339, 346
Speciation, 70–88
Speciation on a conveyor belt: sequential colonization of the Hawaiian islands by *Orsonwelles* spider (Araneae, Linyphiidae), Gustavo Hormiga, Miquel Arnedo, and Rosemary G. Gillespie, 70–88
Species
  boundaries, 159–185
  radiations, 89–109
  trees, 63, 180, 725, 727
*Spermophilus*, 189, 194, 197
  *beldingi*, 189, 204
  *citellus*, 189, 204
  *eversmanni*, 205
  *grammurus*, 204
  *gravojerici*, 204
  *karamani*, 204
  *richardsoni*, 189, 204
  *tridecemlineatus*, 606
  *undulatus*, 189, 204–205
  *variegatus*, 194, 197, 204
*Sphecodogastra*, 24, 32

*Sphenodon punctatus*, 286
Spiders, 56, 58, 60, 62, 70–88, 300, 430, 833, 557
  ground, 430
  *orsonwelles*, 70–88, 833
  *tetragnathid*, 70, 105
Sponges, 525
  asconoid, 313, 332–333
  calcareous, 311, 320, 328, 330, 332–333
  leuconoid, 313, 325, 329
  syconoid, 313, 329, 332
  sylleibid, 313
Spongiae, 520
*Spongilla*, 317
  *lacustris*, 316
Squamata, 159
Squared-change parsimony, 337, 342
Squirrel monkey, 52–53
Squirrels,
  ground, 186, 197
  rock, 194, 197
  terrestrial, 187–188
  tree, 188, 197
Stability, 55–65
  analysis, 554–564
Stability in phylogenetic formulations and its relationship to nodal support, Gonzalo Giribet, 554–564
*Stagonolepis*, 247–248
  *robertsoni*, 241, 243, 250–251
  *wellesi*, 241, 243, 250–251, 254
Stalk-eyed flies, 32
Staphylinidae, 430
Star
  topology analysis, 481
  tree, 236, 236
State (see character state)
Statistical
  analysis, 131, 847
  test, 197
Staurorrhaphidae, 314, 328, 333
*Stenomphrale teutankameni*, 756
*Stephanocemas*, 224
Strepsirrhine, 711–713
Stochastic mapping of morphological characters, John P. Huelsenbeck, Rasmus Nielsen, and Jonathan P. Bollback, 131–158
Stochastic model, 152, 705, 761, 833
Stratiomyidae, 756
Stratiomyomorpha, 746, 750–751, 756
*Straussia*, 820, 826, 828–829, 832–833
Strepsirhini, 738
*Streptopelia*
  *capicola*, 38, 46
  *decaocto*, 38
  *senegalensis*, 38, 46
  *vinacea*, 38, 46
Strict component consensus (SCC), 241, 243, 248, 250
  tree, 251, 532–533, 535
Strict consensus, 406
  tree, 57–58, 61, 323, 339, 344, 737, 826–827
Strict reduced consensus (SRC) method, 240, 243, 247–248, 250
  tree, 250
*Strobilanthes*, 543
*Struthio camelus*, 388, 405
Struthioniformes, 551
*Sturnira lilium*, 472
*Stylipus*, 383
*Suberites*, 317
  *ficus*, 316
Substitution
  model, 3, 140, 293, 371, 463, 577
  rates, 471
Suchard, Marc A., Christina M. R. Kitchen, Janet S. Sinsheimer, and Robert E. Weiss, Hierarchical phylogenetic models for analyzing multipartitie sequence data, 649–664
Suchard, Marc A., Robert E. Weiss, and Janet S. Sinsheimer, Testing a molecular clock without an outgroup: derivations of induced priors on branch-length restrictions in a Bayesian framework, 48–54
*Sudila*, 24, 32
Suine artiodactylans, 854
Sulawesi, 794–819
Sullivan, Jack, *see* Minin, Vladimir, 674–683
Summary characters, 337, 342
Sum of squared deviations (SSD),
Supermatrix, 724–729
  analysis, 724, 726, 728, 853
Supertree, 439–459, 528–538, 724–729, 865
  analysis, 725–726, 728, 853
  carnivore, 845
  clade, 840–841
  consensus, 440, 447
  construction, 728, 839
  MRP, 439, 444
  phylogenetic, 839–851
  support, 55–65, 665–673, 839–851
Supertrees are a necessary not-so-evil: a comment on Gatesy et al., Olaf R. P. Bininda-Emonds, Kate E. Jones, Samantha A. Price, Richard Grenyer, Marcel Cardillo, Michael Habib, Andy Purvis, and John L. Gittleman, 724–729
Supriatna, Jatna, *see* Evans, Ben J., 794–823
Susko, Edward, Chris Field, Christian Blouin, and Andrew J. Roger, Estimation of rates-across-sites distributions in phylogenetic substitution models, 594–603
Svennblad, Bodil, *see* Erixon, Per, 665–674
Swallowtail butterflies, 589
Swofford-Olsen-Waddell-Hillis test (SOWH), 607, 611, 615
Sycetta, 314–315, 332
Sycettidae, 314
*Sycettusa*, 314–315, 320, 322–324, 328, 332–333
  sp., 316
*Sycon*, 311, 313–315, 317, 320, 323, 325, 329, 332
  *calcaravis*, 316–317, 323, 329
  *ciliatum*, 316–317, 322–323, 328–329
  *ensiferum*, 333
  *raphanus*, 316–317, 322–325, 329–330
  spp., 320
*Syconessa*, 314–315, 332–333
Sylvicola alternatus, 756
Symbionts, 735, 743
Symmetric-difference tree, 463
Sympatric
  lineages, 46–476
  speciation, 618–640
*Symphoromyia hirta*, 756
Symphyla, 557
*Synalpheus* shrimp, 631
*Synechocystis*, 659
Syrphidae, 756
Systematic characters, 243
Systematics, 23–36, 55, 259–271, 403, 430, 515–527, 539, 565, 632, 674, 853

molecular, 430, 501–514, 571–572, 591, 745–756, 820–841
phylogenetic, 510
Systematics of the lizard family Pygopodidae with implications for the diversification of Australian temperate biotas, W. Bryan Jennings, Eric R. Pianka, and Stephen Donnellan, 757–780
Systematic utility of floral and vegetative fragrance in two genera of Nyctaginaceae, Rachel A. Levin, Lucinda A. McDade, and Robert A. Raguso, 334–351
Sytsma, Kenneth J., *see* Nepokroeff, Molly, 820–841

Tabanidae, 746, 753–754, 756
Tabanomorpha, 745–746, 750–751, 756
*Tabanus*, 754
*rufograter*, 756
*Taihangia rupestris*, 382
Tamiini, 189, 194
*Tamias*, 189, 205
*sibiricus*, 205
*striatus*, 189, 205
*Tamiasciurus hudsonicus*, 606
Tamura and Nei model, 703
Tandem duplication trees, 110–118
Tapirs, 857
Tarsier, 52–53
Taxic-homology, 539–548
*Taxidea*, 590
*taxus*, 571–573, 581
Taxidiinae, 571–573
Taxonomic
congruence, 853, 862
impediment, 428–435
sampling, 832
training, 428–435
Taxonomic impediment overcome: NSF's Partnerships for Enhancing Expertise in Taxonomy (PEET) as a model, James E. Rodman and Jeannine H. Cody, 428–435
Taxonomy, 330, 390, 474, 539–548
Calyptraeid, 619
Taxon sampling, 71, 119–124, 124–126, 207, 306–307, 423, 442, 446, 451, 539, 542–543, 614, 619, 632–633, 641–642, 644–645, 802, 853
Taxon sampling, bioinformatics, and phylogenomics, Michael S. Rosenberg and Sudhir Kumar, 119–123
Tayra, 573
*Tegula*, 619, 631, 633
Teiidae, 689
*Teinobasis*, 91, 94, 96
*palauensis*, 102
*rufithorax*, 102
Telemetacarpalia, 207, 215, 219
Templeton tests, 15–22, 339, 344–345
Termites, 23, 33
Terrestrial squirrels, 187–188
Tertiary, 549–553, 746, 753
mid-, 549
Testing a molecular clock without an outgroup: derivations of induced priors on branch-length restrictions in a Bayesian framework, Marc A. Suchard, Robert E. Weiss, and Janet S. Sinsheimer, 48–54
Tests
approximately unbiased, 781–793
Bayesian, 607, 615
chi-square, 307, 395, 397–398, 607
goodness-of-fit, 464, 471, 674
hypothesis, 48–54, 781–793
incongruence length difference (ILD) (see also partition homogeneity), 3, 11, 15, 26, 28, 75–76, 418, 467, 604, 607–609, 613, 622, 773, 853–855, 860, 862
Kishino–Hasegawa, 4, 10, 15–22, 212, 216, 390, 392, 394–396, 398, 445, 504, 781
Kolmogrov–Smirnov, 190
likelihood ratio (LRT), 41–42, 48, 54, 63, 95, 352, 363–364, 377, 379, 390, 394, 445, 575, 580, 598, 600, 604–617, 621, 655, 674–683, 714, 801, 825, 849
likelihood weight, 18–19, 22
log-likelihood ratio, 3–4, 377, 379
Mantel, 491
maximum likelihood, 614
Monte Carlo randomization, 684
nonparametric, 781
nonparametric signed–rank, 16
one-tailed, 28, 339
pairwise, 339, 344
parametric bootstrap, 607, 802, 807, 809, 812
parametric, 790
partition homogeneity (see also incongruence length difference), 6, 15, 41–42, 45, 344, 347
phylogenetic, 781
Shimodaira–Hasegawa,15–22, 96, 98, 212, 216, 221, 444–445, 464, 468, 504, 508, 781–793
significance, 781–793
Spearman's rank correlation, 339, 346
statistical, 197
Swofford-Olsen-Waddell-Hillis, 607, 611, 615
Templeton, 15–22, 339, 344–345
topological ILD, 75–77
two-tailed, 339
Wilcoxon rank sum, 337, 345
Wilcoxon signed rank, 406, 668, 857
*Tetilla japonica*, 316
*Tetracerus*, 221
*quadricornis*, 221
Tetraconata, 557
*Tetragnatha*, 103
Tetragnathidae, 56–57
Tetragnathid spiders, 70, 105
*Tetramolopium*, 820
Tetraphyllidea, 430
Tetrapods, 283, 285
evolution,
phylogeny, 283–295
Thamnophiini, 447
*Thamnophis*, 447, 451
*butleri*, 440
*elegans*, 440
Therevidae, 430, 750, 756
Thin-plate spline, 189, 192–193, 488–500
*Thomomys bottae*, 606
*Thoracosaurus*, 398, 401, 417–418
Thorne, Jeffrey L., *see* Wiegmann, Brian M., 745–756
Three-item statements, 255–259
Three-parameter TN92 model, 293
Thrincohalictus, 24
Thrips, 23
Ticks, 296–310
hard, 296–298, 300–301, 304, 309
metastriate, 297, 306
prostriate, 297
rhipicephaline, 304, 306
soft, 300
Tigers, 794

Time flies, a new molecular time-scale for Brachyceran fly evolution without a clock, Brian M. Wiegmann, David K. Yeates, Jeffrey L. Thorne, and Hirohisa Kishino, 745–756
Time-homogeneous model, 647
*Tipula* (*Platytipula*) *paterifera*, 756
Tipulidae, 756
Tipulomorpha, 756
*Tockus nasutus*, 286
*Tomistoma*, 386, 391–392, 394, 396–399
  *cairense*, 405
  *schlegelii*, 386, 388, 403, 405–406, 408–409, 412, 415–419
Tomistominae, 403, 405, 416–418
Topological ILD test, 75–77
Tortricine moths, 430
Total evidence, 55–65, 864
  analysis, 839
Total evidence versus relative evidence: a response to O'Leary et al. (2003), Gavin J. P. Naylor and Dean C. Adams, 864–865
*Trachemys scripta*, 286
Tragelaphini, 209–210
*Tragelaphus*, 209, 212, 219
  *imberbis*, 210
*Tragopan satyra*,
Tragulidae, 206–207, 209–210, 213, 222
Tragulina, 206–207, 209, 218, 222
*Tragulus*, 206, 209, 212–213, 215, 220
  *javanicus*, 210
  *napu*, 210
Transition/transversion
  probabilities, 363
  (TI:TV) ratio, 4, 140, 372, 390, 444–445, 467, 555–556, 559, 577, 650–651, 656, 700, 708–709, 727, 748, 760, 851
Tree-bisect-reconnect (TBR), 120
  branch-swapping, 39, 75, 166, 275, 318, 336, 377–378, 390, 407, 444–445, 463, 504, 506, 575, 607, 621–622, 666, 681, 748, 760, 801, 823–824
Tree-construction analysis, 182
Tree landscapes, 368–373
Tree of Life, 431, 434, 525
Tree squirrel, 188, 197
Trees
  Adams consensus, 463, 850
  allozyme, 509
  Bayesian, 30, 100, 167, 455, 457, 589, 750, 762, 764, 767, 771, 773, 802
  Bayesian consensus, 622
  best-fit, 4
  bifurcating, 642
  binary, 114
  biochemical, 403
  bootstrap, 251, 380–381, 451, 455, 465
  bootstrap consensus, 609
  burn-in, 607
  canonical, 113, 115
  clocklike, 233
  combined data, 612
  consensus, 55, 128, 140, 241, 251, 255, 258, 344, 447, 531, 850, 855
  constrained, 15–18, 20, 22, 761
  constraint, 407, 847, 807
  copaths, 45
  DNA, 770, 773, 775
  equally-parsimonious, 528, 530
  equally weighted, 211
  equal rate, 233
  evolutionary, 526, 602
  gene, 63, 179, 193, 258, 374, 446, 583–584, 588, 725, 727, 776
  genealogical, 515, 517, 521, 524, 526
  haplotype, 159, 165–166
  heuristic, 607
  host, 737, 740
  likelihood, 39, 506, 508, 762, 764, 802
  logDet, 4, 7, 614, 621
  matrix-weighted, 622
  maximum likelihood, 12, 18, 20, 22, 30, 42–45, 81, 96, 128, 323, 328, 368, 371, 383, 396, 455–456, 479, 574, 576–577, 580–581, 583, 589–590, 607, 611, 666, 677, 681, 762, 767, 771, 781, 825, 827–829, 849
  maximum parsimony, 4, 110, 213, 215, 217, 240, 242–243, 248, 252, 275, 317, 320, 323, 329, 336, 339–341, 346, 359, 378, 383–384, 444, 446–447, 463, 574, 576–577, 579–581, 589–590, 607–608, 824, 826–828
  maximum parsimony combined, 16
  minimum evolution, 506, 580, 610
  minimum length, 415, 750
  mitochondrial DNA, 178
  model, 231, 423, 427, 540, 542, 761, 841–842
  molecular, 103, 105–106, 386, 389–392, 394–396, 399, 403, 853, 858, 860
  morphological, 387, 392, 394–396, 409, 503, 860
  most-parsimonius, 3, 5–7, 15, 55–60, 63, 575, 609, 621, 645, 801, 804, 809, 826, 857–857, 864
  multiple, 849–851
  near optimal, 696
  neighbor-joining, 211, 370, 455, 462–463, 508, 702, 801
  nontrivial, 485
  optimal, 4, 8, 17, 20, 121, 128, 320, 341, 357, 696, 760, 850
  parasite, 740, 742
  parsimony, 9–10, 39, 42–43, 79, 286, 645, 762, 764, 771, 773, 802–804
  phylogenetic, 119, 121, 126, 71, 112, 131, 134–135, 244, 289–290, 359, 364–365, 427, 463, 515, 554–555, 669, 671, 685, 692, 715, 749, 760–761, 771, 791, 824, 847, 849
  polyphyletic, 525
  pygopodid, 757, 760, 770, 775
  random, 30, 115, 445, 681
  rodent, 615
  rooted duplication, 110–111, 113, 115
  rooting, 757–780, 865
  single-gene, 20
  source, 727, 840–844, 846
  species, 63, 180, 725, 727
  SSR, 610–611
  star, 236
  strict component consensus, 251
  strict consensus, 57–58, 61, 323, 339, 344, 737, 826–827
  strict reduced consensus, 250
  symmetric-difference, 463
  tandem duplication, 110–118
  unconstrained, 761
  unrooted, 50, 865
  weighted, 866
Triassic, 239–252, 749
  late-, 549, 745, 753
  lower/middle, 746
  mid-, 548
  upper, 746
*Trichogypsia*, 314–315, 317, 323
Trichogypsiidae, 314
Trichomycetes, 430
Trichoptera, 106
*Trichotropidae*, 639
  *cancellata*, 639
  *conica*, 639
  *insignis*, 639

Trichotropids, 620, 623
Trigonioida, 559
*Trimeresurus*, 447, 451
  *albolabris*, 441
  *tokarensis*, 441
*Tripedalia cystophora*, 316
Triticeae, 383
tRNA, 2, 52, 94–95, 139, 296–298, 300–301, 303–308, 383, 396, 406, 460, 615, 725, 727, 758, 762, 798
  sequences, 296, 298
tRNAAsp, 674
tRNAL, 73, 76
*trn*L-*trn*F, 380–382, 384
*Trochita*, 619, 623, 631, 639
  *calyptraeformis*, 631, 639
*Trogon collaris*, 286
Tropidophiidae, 440
*Tropidophis*
  *haitianus*, 440
  *wrighti*, 440
*Tropidurus*, 689
True and false gharials: a nuclear gene phylogeny of Crocodylia, John Harshman, Christopher J. Huddleston, Jonathan P. Bollback, Thomas J. Parsons, and Michael J. Braun, 386–402
True sampling, 665–673
Tuatara, 286
*Tulearinia*, 329
Tunicata, 520
*Turdus migratorius*, 287
*Turitella* sp., 640
Turkey, 551
*Turnix sylvatica*, 286
*Turtur brehmeri*, 38
Two-state characters, 236
Two-tailed test, 339
Tylopoda, 221
Type I error, 18, 42, 273, 279, 505, 615, 666–669, 671–672, 684, 689
Type I STS markers are more informative than cytochrome *b* in phylogenetic reconstruction of the Mustelidae (Mammalia: Carnivora), Klaus-Peter Koepfli and Robert K. Wayne, 571–593
Type II error, 273, 279, 505, 666, 671–672, 770
*Typhlomolge*, 504
Typhlopidae, 440
*Typhlops lumbricalis*, 440
*Typhlotriton spelaeus*, 505
*Typothorax*, 241, 243, 248–252
  *coccinarium*, 254
*Tyrannus tyrannus*, 286

U2 snRNA, 557
Ulocentra, 492–493
Unambiguous characters, 539, 557
Unbiased (see approximately unbiased test)
Unconstrained
  ordination, 687
  trees, 761
Uninformative characters, 76, 217, 575, 581, 826
Unionoida, 559
Ungulates, 860
Unrooted tree, 50, 865
Unweighted
  pair-group average (UPGMA), 205, 229–239, 722
UPGMA
  cluster analysis, 488, 491, 493, 496–497
  phenogram, 497
*Upupa epops*, 286
*Urocolius macrourus*, 286
Uropygi, 557
Ursidae, 846
*Ursus*, 706
Using a null model to recognize significant co-occurrence prior to identifying candidate areas of endemism, Austin R. Mast and Reto Nyffeler, 271–280
*Ute*, 314–315
*Uteopsis*, 314–315

Vacelet, Jean, *see* Manuel, Michael, 311–333
*Vallesia*, 391,
Value of idiosyncratic markers and changes to conserved tRNA sequences from the mitochondrial genome of hard ticks (Acari: Ixodida: Ixodidae) for phylogenetic inference, Anna Murrell, Nick J. H. Campbell, and Stephen C. Barker, 296–310
*Vanikoro*, 622, 639
  sp. 1, 639
  sp. 2, 639
Varanidae, 757
*Varecia*, 706
Variable rate analysis, 751
Vermileonidae, 756
Vertebral characters, 439
Vertebrate, 56, 58–59, 261–262, 264, 296, 388, 403, 515, 518, 524, 526, 571, 706
  phylogenetics, 388
Vespid wasps, 23
Vicariance, 549–553, 717–723, 794–819
Villet, Martin H., *see* Kelly, Christopher M., 439–459
Violet turaco, 388
*Vipera*, 451
  *ammodytes*, 441
  *ursinii*, 441
Viperidae, 439, 441, 447, 452–453, 455
Viperinae, 441, 447, 453
Viverridae, 846
*v-myc*, 388
Volatiles, 334–351
*Vombatus ursinus*, 285
Vos, R. A., Accelerated likelihood surface exploration: the likelihood ratchet, 368–373
*Vosmaeropsis*, 314–315, 320, 322–325, 328, 333
  sp., 316
Vulture, black, 388
  vWF, 605, 614–615, 738

Wagner, Warren L., *see* Nepokroeff, Molly, 820–841
*Waldsteinia*, 374, 381–383
  *geoides*, 374, 376, 381–382
Wallace's Line, 794–819
Wasps
  parasitoid, 430
  vespid, 23
Watson–Crick pairing, 131
*Waxy* (see also Granule-bound starch synthase), 374–385
Wayne, Robert K., *see* Koepfli Klaus-Peter, 571–593
Weasels, 571–573
  long-tailed, 573
Weevils, 717
Weighted
  parsimony, 283
  tree, 866
Weiss, Robert E., *see* Suchard, Marc A., 48–54, 649–664
Whales, 707–708, 724, 853, 855, 858
  baleen, 707–708

fin, 210
phylogeny, 853
toothed, 707–708
When are phylogenetic analyses misled by convergence? a case study in Texas cave salamanders, John J. Wiens, Paul T. Chippindale, and David M. Hillis, 501–514
When do parasites fail to speciate in response to host speciation?, Kevin P. Johnson, Richard J. Adams, Roderic D. M. Page, and Dale H. Clayton, 37–47
Wiegmann, Brian M., David K. Yeates, Jeffrey L. Thorne, and Hirohisa Kishino, Time flies, a new molecular time–scale for Brachyceran fly evolution without a clock, 745–756
Wiens, John J., Missing data, incomplete taxa, and phylogenetic accuracy, 528–538
Wiens, John J., Paul T. Chippindale, and David M. Hillis, When are phylogenetic analyses misled by convergence? a case study in Texas cave salamanders, 501–514
Wilcoxon
rank sum, 337, 345
signed rank test, 406, 668, 857
*T*-distribution, 345
Wilkinson, Mark, François-Joseph Lapointe, and David J. Gower, Branch lengths and support, 127–130
Wilkinson, Mark, *see* François-Joseph Lapointe, 865–868
Wilkinson, Mark, *see* Harris, Simon R., 239–252, 851–852
Williams, David M. and Christopher J. Humphries, Component coding, three-item coding, and consensus methods, 255–258
Winka, Katarina, *see* Cummings, Michael P., 477–487
Winteraceae, 346
Wishart distribution, 652
Wolverine, 573
Woodchuck, 193, 195, 197–198
Wombat, 285
Wright–Fisher model, 179

Xenoderminae, 439, 441, 447, 452–453, 457
*Xenodermus*, 451
*javanicus*, 441
Xenodon, 447, 451
*neuwiedi*, 441
*servus*, 441
Xenodontinae, 439, 441, 447, 452, 455–456
Xenodontini, 447, 456
*Xenophora pallidula*, 640
*Xenopus laevis*, 286
*Xenopus laevis*, 286
Xia, Xuhua, Zheng Xie, and Karl M. Kjer, 18S Ribosomal RNA and tetrapod phylogeny, 283–295
Xie, Zheng, *see* Xia, Xuhua, 283–295
Xiphosura, 557
*Xylocopa*, 8
*bombylans*, 2, 5
Xylocopinae, 1–2, 9
Xylocopini, 1, 3, 9
*Xylomya parens*, 756
Xylomyidae, 746, 756
Xylophagidae, 756
Xylophagid flies, 745
Xylophagomorpha, 745–746, 750–751, 756

Yacare caiman, 388
Yang, Ziheng and Anne D. Yoder, Comparison of likelihood and Bayesian methods for estimating divergence times using multiple gene loci and calibration points, with application to a radiation of cute-looking mouse lemur species, 705–716
Yeates, David K., *see* Wiegmann, Brian M., 745–756
Yoder, Anne D., *see* Yang, Ziheng, 705–716

Zapodidae, 614
*Zapus*, 614
*princeps*, 606
*Zeacrypta*, 636
*Zegalerus*, 632, 639
*tenuis*, 639
*Zenaida*, 45
*galapagoensis*, 38, 45
Zimmer, Elizabeth A., *see* Nepokroeff, Molly, 820–841
Zorilla, 573
Zwickl, Derrick J., *see* Hillis, David M., 124–126
Zygethobiini, 559
*Zygethobius*, 559